

THE PLATE CLIMATOLOGY THEORY

How Geological Forces Influence , Alter, or Control Earth's Climate and Climate Related Events

Written by: James Edward Kamis

plateclimatology.com • plateclimatology@gmail.com

ACKNOWLEDGEMENTS

Theory technical consultation provided by; meteorologist and computer programmer Paul E. Kamis, geophysicist James Ligon, geologist Steven Parks, and geologist David Entzminger. Thanks to Steve Parks and James Ligon for final editing and years of lively discussion concerning many aspects of the theory. Thanks to Laura Wilson for building and maintenance of the theory website. Thanks to Boston free-lance writer Thomas Richard and the editor of the Climate Change Dispatch website for posting my many climate articles. Special thanks to my wife Nadine Ellen Schoenbeck Kamis for moral support and encouragement during the lengthy process of developing this theory.

ABSTRACT

Increased tectonic and volcanic activity, either locally or globally, equates to more geologically induced heat transfer and chemically charged temperature altered fluid (HCF) release from active geological features into oceans, polar areas, and atmosphere. Releases, that have in past and still to this day act to significantly alter or completely control earth's climate and climate related events. To describe this new theory, the term Plate Climatology was designated on 7 October 2014 ([Kamis 2014](#)) and presented at the 2016 annual American Metrological Society Conference ([Kamis 2016](#)).

The significant climate effect of geological forces has been hidden from scientific investigation and understanding for several reasons. First, most of the HCF is from three extensive, remote, under explored, and under monitored regions. Specifically, the Antarctic, Arctic and deep oceans. These regions represent; 82% of earth's surface, most of earth's outer crustal plate boundaries, and numerous high HCF volcanic complexes. Virtually none of these geological features are monitored for HCF emission. Secondly, the modern-day

worldwide volume and percent of geologically induced CO2 and methane emissions from terrestrial regions has been greatly underestimated. Third, the limited amount of HCF data from remote geological features has led to the incorrect assumption that these features do not significantly contribute to changes in climate. Lastly, failure to appreciate and then incorporate into current climate theory the significant influence of ancient geologically induced extinctions, ice ages, ocean anoxia, ocean current alterations, and El Nino / La Nina events.

DISCUSSION

The Plate Climatology Theory provides a powerful platform to evaluate, verify, and integrate; previous geological postulations, relevant information, and the author's observations. This process has allowed the author to achieve a more accurate realization of the significant climate impact of geological forces. In a very broad sense the main premise of the Plate Climatology Theory, that geological forces significantly influence climate, is substantiated by appreciating:

- The immense power of earth's largest geological features, upper mantle convection systems. Systems that have the energy to move entire continents 2.5 centimeters per year by emitting massive amounts of lava and associated chemicals, minerals, gases, water, and heat into earth's outer crust, oceans, and atmosphere. Knowing this it is reasonable to theorize that upper mantle convection cells have the power to significantly effect earth's climate and climate related events.
- The numerous examples of locally juxtaposed; melting to non-melting glacial ice areas, melting to non-melting ice sheets, warm to cool ocean areas, and warm to cool atmospheric areas. The boundaries between these local areas are distinct, well defined, and often of significant magnitude. It is difficult to reconcile all of these boundaries with forces associated with non-local, uniform, and worldwide atmospheric global warming. A better explanation is that many, but not all, are the result of HCF from geographically distinct geological features located in the boundary areas.
- The consistent failure of atmospheric based computer climate models to accurately predict future changes in climate trends or to reliably define the root cause of climate current trends. Failure of these climate models is likely the result of not properly including the effect of geologically induced HCF.
- That ancient extinction events, ocean anoxic events, beginnings of glacial periods, and endings of glacial periods are proven or very likely to be the result of geological forces.

The basic principles of the Plate Climatology Theory were conceived in the late nineteen seventies by considering the importance and interrelationship of two significant advances in science. First, the discovery of an entirely new environmental system located on our ocean seafloors termed the Chemosynthetic System. An environmental system that is fueled by massive geologically induced HCF emissions and unrelated light from the sun. Secondly, verification of Alfred Wegener's Continental Drift Theory which stated that inner earth geological forces had the power to move entire continents. It occurred to the author that if geological forces had the power to move continents and support a huge environmental system, they certainly had the power to influence earth's oceans, atmosphere, and climate.

The significant climate effect of geological forces has remained hidden and underappreciated by scientific investigators because their heat and fluid release are primarily found in under explored, remote, or under monitored regions:

Polar Ice Caps are covered by an average ice thickness of 1,524 meters (5,000 feet), account for 11% of earth's surface, and are home to major geological features such as: divergent plate boundaries, transform plate boundaries, and high heat flow volcanic regions.

Oceans which account for 71% of earth's surface, have an average depth of 4,267 meters (14,000 feet), and contain most of earth's major geological features such as divergent plate boundaries, transform plate boundaries, convergent plate boundaries, and high heat flow volcanic regions.

Terrestrial Volcanic Regions are poorly understood because very few of their estimated 1,500 volcanoes are monitored for emissions of CO₂, methane, sulfur, hydrogen, mercury, chloride, particulate matter, and heat. New research has shown that non-erupting volcanoes and associated volcanic features can contribute significantly to the CO₂ and methane content of our atmosphere ([Ilyinskaya 2018](#))

Even after years of research the rate, magnitude, volume, and chemical composition of emissions from geological features are still poorly understood. These emissions include heat transfer, liquid fluid flow (primarily heated seawater, cooled seawater, CO₂, methane, sulphur, iron, phosphorous, and mercury), and gaseous fluid flow (primarily heated air, CO₂, and methane). To describe these geologically induced emissions the term "heat flow and chemically charged temperature altered fluid flow" (HCF) is referred to throughout this paper.

The idea that geological forces influence earth's modern and ancient climate has been postulated by numerous scientists for many years. The author has attempted to properly credit all these scientists and their ideas in the text or reference sections of this paper.

Lastly, the Plate Climatology Theory is not intended to be a replacement of all aspects of current climate theory, rather a plausible alternative explanation of selective; climate events and trends, climate related events and trends, and individual aspects of current climate theory.

EVIDENCE

Polar Ice Caps

The end of Glacial Periods signified by retreat of glacial ice sheets begins suddenly and transpires in a short time frame relative to the gradual beginning and long duration of Glacial Periods represented by ice sheet advances ([Steffansen 2008](#)), [Blatter 2013](#), [Abe-Ouchi 2013](#), [video simulation Abe-Ouchi 2013](#), and [Van Ommen 2011](#)). The author theorizes that these rapid Glacial Period endings are the result of worldwide pulses of increased volcanism from both ocean floor ([Martínez-Botí 2105](#)) and terrestrial volcanic ([McConnell 2017](#)) geological features.

Ocean floor geological features act to warm and chemically alter seawater. Seawater warming is the result of geothermic heat transfer and emissions of superheated water from geological features such as; ocean floor hydrothermal vents, seamounts, active faults, volcanic complexes, and spreading centers. Emissions of chemically charged fluid, primarily CO₂ and methane, from these same geological features acts to alter the chemistry of seawater. Alteration of seawater also occurs when geologically warmed oceans decrease their CO₂ concentration.

Chemical alteration of the atmosphere is the result of greenhouse gas releases from the underlying oceans and terrestrial sub-glacial volcanic features. Atmospheric warming is the result of increased CO₂ concentrations, increased methane concentrations, and heat transfer from warmed oceans.

Glacial Periods are known to begin suddenly but last for long periods of time relative to Glacial Period endings. ([Abe-Ouchi 2013](#), [Steffansen 2008](#)). The author theorizes that the sudden cessation of a worldwide HCF pulse marks the end of Polar Ice Sheet retreats and the beginning of a so-called Glacial Period. Advance and resulting

increased geographical extent of Polar Ice Sheets is not the result of anomalous forces, rather a recovery to the normal geographical extent of earth's ice sheets. As a result, the term Glacial Period does not accurately describe what is actually a very long and natural recovery process. The term Glacial Period should be replaced with the term Glacial Recovery Period.

Evidence supporting this new explanation for Glacial Period processes comes from several sources. First, the close time correlation of beginnings and endings of significant worldwide volcanic pulses with the beginnings and endings of Glacial Periods ([Jagoutz 2016](#) and [Muschiello 2017](#)). Secondly, geological history provides numerous analog recovery processes following significant HCF pulses for example large extinction ([Jerram 2016](#)) and ocean anoxic events ([Jenkyns 2010](#)). These HCF induced events typically occur suddenly and within a relatively short time period. In contrast, the recovery process typically takes significantly more time. Another analog is the alteration of coral reefs by El Nino ocean temperature increases and chemical alterations. El Nino a natural rapidly occurring ocean HCF event. Reef recovery is complex and longer term.

Failure of many computer models to accurately simulate or explain the beginnings and endings of glacial periods is because these models utilize input data that are side effect data of the worldwide HCF pulses. These side effect data include but are not limited to changes in wind, atmosphere temperatures, ocean currents, ocean temperatures, sedimentation sequences, etc. Researchers have in past written many papers attempting to explain the root cause of Glacial and Inter-Glacial Periods. These explanations are often complex, tedious, and unconvincing likely because they invoke data from side effects HCF pulses.

The sudden onset of rapid retreat and melting of selected portions of our polar ice caps mimics past HCF induced endings of Glacial Periods as described above. This is strong evidence supporting the idea that HCF and not anthropogenic warming is the primary cause of changes to earth's polar ice caps.

Antarctica

Portions of Antarctica's glacial ice sheet, sea ice, surrounding ocean waters, and biological regimes are currently undergoing significant change. Those advocating the theory of Global Warming / Climate Change have championed these changes as proof positive that human CO2 emissions are at work in Antarctica. Implicit in their argument is that the underlying bedrock geology of Antarctic is of little or no consequence and merely acting as a passive support system for seawater and glacial ice. However, numerous research studies, especially those released within the last ten years, demonstrate that many of Antarctica's bedrock geological features are not passive, instead extremely active. Furthermore, the research proves that these active geological features are emitting massive amounts of HCF into Antarctic oceans and bedrock ice interfaces, thereby acting to change Antarctica's glacial ice sheet, sea ice, surrounding ocean waters, and biological regimes. A more detailed discussion of evidence supporting this contention is as follows.

The Antarctic continent which is greater in areal extent than the lower 48 US states, is formally divided into two land mass segments; West Antarctica 20% (note that the designation "West Antarctica" includes the West Antarctic Peninsula) and East Antarctica 80%. Recent NASA research, as per the Figure 1 graph, demonstrates that 100% of the Antarctica Continent's ice sheet loss, here expressed as global sea level rise per year, can be attributed to the much smaller West Antarctica region. The graph also shows that East Antarctica is gaining rather than losing ice.




Figure 1.) Map shows red outlined West and East Antarctic land masses. Overlying graph illustrates Antarctic Ice Sheet contribution to global sea level rise by geographic region which is a proxy for ice loss ([Ivins, E., 2018](#)).

Other NASA research shows that the ice sheet gains in East Antarctica outpace ice sheet loses in West Antarctica and as a result the Antarctic Continent is gaining, not losing ice ([Zwally 2015](#)). So, what force or forces can explain ice gain across 80% of the Antarctic Continent while at the same time the immediately adjacent 20% is losing ice?

The answer becomes evident by reviewing Figure 2 which is the University of Washington's Antarctic Fifty-Year Average Surface Temperature Map ([Steig 2009](#), and [Nicolas 2014](#)). This map demonstrates that there are two distinct and very different near surface temperature areas. One area of anomalously high near surface temperatures associated with West Antarctica and another area of low near surface temperatures associated with East Antarctica.

The author theorizes that the high surface temperatures in West Antarctica are the result of HCF emissions from numerous geological features associated with Antarctica's largest and most active geological feature, the West Antarctic Rift System. Absence of widespread or significant HCF in the East Antarctic area accounts for near average near surface temperatures in this portion of Antarctica.




Figure 2.) Antarctic fifty-year average Surface Temperature Map. Areas of normal and high surface temperatures interpreted by author as high and low heat flow areas (research credit Steig, E., 2009, heat flow region outlines credit [Kamis, J. E., 2016](#)).

The validity of Dr. Steig's Figure 2 map was hotly debated for many years, however it has since been proven correct by two more modern research studies, NASA's Skin Temperature Map ([NASA, 2006](#)) and the British Antarctic Survey's Geothermal Heat Flux Map ([Martos, Y., et. al., 2017](#)). Dr. Steig's map was considered controversial because it did not demonstrate, as per Global Warming principles, an evenly distributed continent-wide near ground surface warming pattern. An even widespread pattern could have been interpreted as evidence uniform manmade atmospheric global warming was at work in Antarctica by melting the top surface of the ice sheet. Scientists now widely accept that loss of Antarctic Ice Sheet mass and extent is the result of basal, not top melting which can be interpreted as the result of bedrock HCF ([Kamis 2018](#), [NASA 2017](#), and [Jordan 2018](#)).

Other continent-wide information indicating that many modern-day changes in Antarctica are related to geologically induced HCF and not manmade atmospheric global warming are:

- The Overall East Antarctica ice gains have outpaced West Antarctica ice losses, as a result the Antarctic Continent has been gaining, not losing ice mass continuously for many years ([Zwally, J., 2015](#)).
- The atmosphere above the Antarctic Continent has been cooling, not warming for many years ([Sejas, S. A., 2018](#), [Abrams, N. 2014](#), [Turner, J., et. al., 2016](#), [Hinkel, L., 2016](#)).
- The ocean waters immediately adjacent to West Antarctica are dramatically warming while ocean waters adjacent to East Antarctica are only mildly warming or not warming at all. ([NASA, April 2006](#)).
- The existence of an extensive subglacial interconnected system of liquid freshwater lakes and streams, most significantly above sea level. ([Wright, A., et. al., 2012](#), [Fricker 2007](#), [Siegert 2016](#), [Goller 2014](#)).
- The base and not the top of the West Antarctic Ice Sheet is melting ([Seroussi, H., et. al., 2017](#))

Next, we discuss specific examples of how active geological features effect Antarctica.

[West Antarctic Rift System](#)

The West Antarctic Rift System (WARS) is a 6,437-kilometer-long (4,000 miles) and 1,127-kilometer-wide (700 mile) geological feature that has a long and complicated history of plate boundary subduction, transform movement, and rifting (Figure 3). The intent of this article segment is not to describe the geological history or complexities of this huge rift system, rather to describe the significant effect of its present day and ancient HCF. The WARS is home by latest count to; approximately 138 (De Vries 2018) land volcanoes, a 1,605,793 square kilometer (620,000 square mile) mantle plume, thousands of deep inner earth reaching faults, numerous seamounts, and a countless number of hydrothermal vents. Recent research studies by NASA (NASA 2017), the University of Texas (Schroeder 2014), and many other organizations confirm that large amounts of HCF is currently being emitted from the rift's thousands of active geological features into Antarctica's oceans and atmosphere. These emissions and other geological evidence support the idea that the WARS is a very geologically active part of the Pacific Ring of Fire (Kamis, 2018). Other data supports the idea that the WARS is an active part of the Pacific Ring of Fire:

- Contains a large number of active volcanoes, on a per square kilometer basis the same distribution as the Pacific Ring of Fire, one every 24,000 square kilometers (9,267 square mile) (J. E. Kamis calculation).
- Is home to a large number of active faults, here estimated to be a similar per square kilometer density as the Pacific Ring of Fire.
- Connects in a geologically seamless fashion onto the two currently defined horseshoe ends of the Pacific Ring of Fire located in South America and New Zealand (Kamis 2018, Gurion 2018).
- Has become increasingly active in coincidence with increasing tectonism along the Pacific Ring of Fire, therefore indicating a geological processes connection to the Ring of Fire (Kamis 2018).




Figure 3.) Location of Antarctica's high heat flow and chemically charged high fluid flow major geological features. High bedrock heat flow cross-hatched black and very high bedrock heat flow areas cross-hatched red (credit Google Earth and J. E. Kamis).

[South Georgia and South Sandwich Islands](#)

The South Georgia and South Sandwich volcanic island chains and associated volcanic plateaus were generated by subduction of the South American Plate beneath the Antarctic Plate (Figures 3 & 4, [Barker 1981](#)). These island chains are a geologically active 1,600-kilometer (994 mile) extension of the West Antarctic Rift System originating at a location where the rift's major fault traces conjoin to the northern extension of the West Antarctic Peninsula. The island chain's subduction zone fault system continues westward and northward eventually seamlessly connecting to South America's west side subduction zone. These island chains are currently very geologically active as exemplified by recent multiple earthquakes and volcanic eruptions. On December 10, 2018 there was a 7.3 magnitude earthquake located in proximity to Montagu island of the South Sandwich Island chain. On September 29, 2016 the simultaneous eruption of three South Georgia Island volcanoes was captured by NASA satellites. On May 1, 2016 the South Sandwich Island's Mount Curry eruption spewed ash into the region's atmosphere, ocean waters, and onto ice surfaces. This ash adversely affected the South Sandwich Islands one million resident penguins.

These eruptions and the powerful earthquake can be interpreted as evidence that hundreds of other geological features located along this 1,600-kilometer-long (994 mile) extension of the West Antarctic Rift System may also be actively emitting HCF into adjacent ocean waters, land surfaces, and atmosphere.

Recent research in the South Georgia and South Sandwich Islands area discovered the presence of mercury and methyl mercury in the region's sea ice and ocean waters ([Rodgers 2012](#) and [Gionfriddo 2016](#)). These author's theorized that this mercury and methyl-mercury was human sourced. However, mercury is a common component of terrestrial volcanic ash. Additionally, mercury and methyl mercury are common components of seamount and hydrothermal vent emissions. These types of geological features are abundantly present and very actively emitting large volumes of HCF enriched in iron, mercury, and methyl mercury into in the South Georgia / South Sandwich area ([Klar 2017](#)). Geologically induced emissions are the likely source of the area's mercury and methyl mercury thereby strengthening the notion that geological forces have a wide-ranging and likely dominant effect on this region's climate and climate related events.




Figure 4.) Detailed location map of the northern extension of the West Antarctic Rift System. Rift faults as black lines, active volcanoes as larger yellow filled red circles, and semi-active volcanos as smaller yellow filled red circles. (map credit Google Earth, labels and faults credit J. E. Kamis)

[South Shetland Islands](#)

The 725-kilometer-long (450 mile) and 80-kilometer-wide (50 mile) South Shetland volcanic island chain is an active part of the West Antarctic Rift System. It is home to ten dormant and one currently active volcano, all fueled by West Antarctic Rift System faults (Figure 4). This volcanic chain, and areas immediately adjacent to it, have in past and still to this day emit high amounts of HCF into the overlying ocean thereby acting to warm these ocean water areas. Examples are as follows:

The Deception Island Collapse Caldera which has in past and is still today a very active geological feature. Caldera bay floor hydrothermal vents are currently emitting large amounts of HCF into the caldera's 4 kilometer wide (2.5 mile) circular ocean bay. Emission of these hot fluids into the bay acts to keep much of the bay ice-free and in selected areas elevate the bay water to sauna like temperatures. The Deception Island volcano has erupted ten times from 1827 to 1995. Just released research shows that this caldera violently erupted 3,980 years ago likely acting to change earth's climate for up to twelve months ([Antoniades 2018](#)). The current magnitude and geographical extent of ocean warming and chemical charging by the Deception Island Caldera and its associated geological features is unknown, because like most of Antarctica's geological features it is not emission monitored. However, they are likely significant.

Parallel to, and just east of the South Shetland Island chain is the 450-kilometer-long (280 mile) and 50 kilometer-wide (31 mile) Bransfield Strait. Numerous research studies have concluded that this highly complex geological feature is currently very active ([Barker 1994](#) and [Klinkhammer 2001](#)). The Klinkhammer research states that... "*data on the high heat flow, active volcanism, and extensional folding, in combination with modeling of the gravity and magnetic anomalies and earthquake focal mechanisms, indicate that the Bransfield strait floor represents a zone of lithosphere extension forming in the Antarctic Peninsula. The most important structural element of the strait floor is the neo-volcanic zone near the South Shetland Islands and the diapirism zone near the Antarctic Peninsula.*".

[Larsen Volcanic Plateau](#)

The Larsen Volcanic Plateau is an 81-kilometer-long (50 mile) and 16-kilometer-wide (10 mile) part of the WARS (Figure 3). This plateau and the surrounding sea ice fields have been the focus of many research studies and media articles that conclude that the areas sea ice breakup and melting is the result of manmade atmospheric global warming. An alternative explanation is related to geologically induced HCF from Larsen Volcanic Plateau geological features.

The plateau is home to 16 semi-active / dormant volcanoes and one actively fluid emitting seamount. Volcanoes form prominent circular topographic highs in the ice sheet. Many are void of snow and ice which is a direct indication of ice melting geothermal heat flow.

The breakup of the sea ice adjacent to the Plateau occurs progressively through time farther and farther away from the plateau. This is interpreted to be the result of progressive lateral geologically induced warming of ocean waters in the plateau region. Importantly, other distant areas of Weddell Sea Ice are not melting, just the sea ice adjacent to the Larsen Volcanic Plateau. This indicates that the powerful ice melting heat source is local and not widespread.

Information obtained from the Smithsonian Global Volcanism Program Stated the following;

"In January 1982, geologists from the Universidad de Chile found evidence of recent volcanism at two sites in the Seal Nunataks Group that had not previously been identified as recent volcanic centers. At Dallman (65.02°S, 60.32°W), very fresh-looking basaltic lava had emerged from the central crater and flowed to the NW foot of the volcano. Active fumaroles were observed at the N side of the summit. Abundant basaltic lapilli and ash covered wide areas of the Larsen Ice Shelf in

the vicinity of Murdoch Volcano (65.03°S, 60.03°W). Portions of these deposits were covered by fresh snow. Fumaroles were active in a small parasitic cinder cone SE of Murdoch. In December 1893, C.A. Larsen saw an eruption of black ash from Lindenberg (64.92°S, 59.70°W) and solfataric activity at Christensen (65.10°S, 59.57°W), but no activity was observed at either of these volcanoes in 1982."

Earliest explorers of Antarctica mentioned in their journals the presence of fresh volcanic ash on glacial ice in the Larsen Volcanic Plateau area. One other recent visit by researchers to the ocean seafloor of the Larsen Volcanic Plateau discovered an actively heat emitting seamount ([Dormack 2004](#)). Research by a multi-national team concluded that very rapid uplift in the Larsen Volcanic Complex area in 2002 is difficult to associate with overlying ice sheet melting and subsequent glacial rebound ([Nield 2014](#)). Although not proven, the bedrock uplift event is more likely related to upward movement of deep earth magma as is currently occurring at the Marie Byrd Mantle Plume region along the WARS.

Evidence indicates that the Larsen Volcanic Plateau is a geologically active area which is emitting significant amounts of HCF.

[Hudson Mountain Volcanic Complex, Pine Island Glacial Valley , and Mount Takahe Volcanic Complex](#)

Melting, fracturing, and breakup of glaciers in this area had for many years been attributed to the effects of anthropogenic warming as per the Climate Change Theory. However, more recent research ([Loose 2018](#), [McConnell 2017](#), [Schroeder 2014](#)) combined with the authors observations indicate that modern day and ancient changes of these glaciers are the result of HCF from the areas active geological features, and not anthropogenic atmospheric warming.




Figure 5.) Very High heat flow Hudson Mountains and Mount Takahe Volcanic Complexes (red cross-hatched). Down faulted valleys (black outline). Active subglacial volcanoes (red outline with yellow fill) and active surface volcanoes (red circle). (credit Google Earth and J. E. Kamis)

Location of the Hudson Mountain Volcanic complex, the Mount Takahē Volcanic complex, and the Pine Island Glacial Valley are illustrated on Figure 5. These geological features were created and are fueled by forces associated with the western bounding regions of the WARS ([Kamis 2017](#) and [Kamis 2017](#)). This area's present-day high HCF has been confirmed by numerous research studies ([Dziadek 2018](#), [Steig 2009](#), [Lough 2013](#), [Iverson 2017](#)).

The long linear shape and orientation of the Pine Island Glacial Valley is here thought to be the result of down faulting and graben generation associated with WARS. In June of 2018 a multi-national team of researchers made a stunning discovery. They confirmed the presence of a currently erupting volcano beneath the Pine Island Glacier ([Loose 2018](#)) thereby strongly indicating that eruptive heat flow is melting this valley's glaciers, not Climate Change. Other recent research shows that there are also active volcanoes beneath the adjacent Thwaites Glacier (see Figure 6 and [Kamis 2019](#)). Concerning the Hudson Mountain Volcanic Area, research by the British Antarctic Survey proved that 2,200 years ago a massive sub-glacial volcanic eruption covered this ancient glacial ice area with a thick layer of volcanic ash. Post eruption glacial ice accumulations buried the ash layer ([Corr 2008](#)). Based on the extent and thickness of the ash layer researchers described this event as one of the largest Antarctic eruptions in 10,000 years. Research by Nevada's DR Institute showed that 17,700 years ago Mount Takahē erupted on multiple occasions during a 192-year span ([McConnell 2017](#)). This world class eruptive phase acted to alter ocean currents, ice extent, and climate of the entire southern hemisphere in an area extending from the South Pole to the sub-tropics.

[Marie Byrd Mantle Plume Area](#)

A recent NASA study confirmed what geologists have suspected for many years, that a very large and active mantle plume hotspot is present beneath the glacial ice of West Antarctica (Figure 2 and [NASA 2017](#)). Termed the Marie Byrd Mantle Plume, a subglacial hotspot which fuels and is home to approximately 100 active, erupting, or dormant volcanoes ([De Vries 2018](#)). Several of these volcanoes are currently erupting beneath the glacial ice ([Lough 2013](#), [Schroeder 2014](#), and [Loose 2018](#)). A very recent research study showed that the bedrock surface expression of the mantle plume is uplifting at a very rapid rate ([Barletta 2017](#)). An uplift rate which is greater than can be accounted for by glacial isostatic rebound and is therefore almost certainly related to upward movement of the underlying mantle plume magma chambers.




Figure 6.) Marie Byrd Mantle Plume, Mount Erebus, and Pluton areas (red hatched) in relationship to ice sheet melting (red shading). Black cross-hatched is West Antarctic Rift System west and east branches (credit NASA and J. E. Kamis).

Figure 6 illustrates West Antarctic Ice Sheet melting from 1992 to present displayed as varying red shades. Superimposed on this map are the red hatched outlines of the; 1,605,793 square kilometers (620,000 square mile) of the Marie Byrd Bedrock Mantle Plume "Hotspot" ([Kamis 2018](#)), Mount Erebus Area ([Kamis 2016](#)), 110,000 square kilometer (4,2471 square mile) Pluton Area ([Allibone 2010](#)), and the major bounding faults of the WARS ([Kamis 2016](#)). The outline of these three subglacial geological features closely matches the outline of the most prominent ice sheet melt regions. This correlation is strong evidence that HCF from these subglacial geological features is the dominant force acting to melt the overlying ice sheet.

HCF induced basal melting of the West Antarctica ice sheet in the Marie Byrd Mantle Plume area acts generate a significant amount of warm liquid water. This basal meltwater outflows downhill into adjacent ocean regions where it then acts to significantly warm the outflow seawater area as illustrated on NASA's Antarctica Skin Temperature Map ([NASA 2016](#)). The NASA map also illustrates two other very telling aspects of the ocean waters surrounding the Antarctic Continent. First, the other significantly warmed ocean areas adjacent to West Antarctica are all associated with active geological features. Secondly, East Antarctica which is geologically inactive has only a few adjacent slightly ocean seawater warmed areas. These observations are theorized to demonstrate that warming of ocean waters adjacent to the Antarctic continent are the result of geologically induced HCF, not manmade atmospheric warming.

[Mount Erebus Volcanic Complex / Pluton Rich Area](#)

The Mount Erebus Volcanic Complex is 105,000 square kilometer (40,541 square mile) region that includes three stratovolcanoes, the 5,000 square kilometer (1,931 square mile) McMurdo Dry Valley area, and a portion of an adjacent ocean bay seafloor. All these components lie directly atop and are fueled by an active segment of the WARS (Figure 7).

Mount Erebus has been volcanically active for thousands of years. Records taken from several sources show that eruptions occurred in; 8050 BC, 7050 BC, 4050 BC, 4550 BC, 2950 BC, 2050 BC, 841,1900, 1903,1908, 1911, 1912, 1915,1915,1947, 1955, 1957,1963,1984, 1993, 1995, 1997,1998, 2000, 2001, 2005, 2014, and

2015 ([Volcano Discovery](#) and [Smithsonian Institution](#)). At present the Mount Erebus summit crater is actively pulsing molten lava and gases. Numerous vents located along the flanks of the volcanic cone are also currently emitting heated gases. Research by New Mexico Tech shows that as of 2008 the main magma chamber, which was estimated to be one half mile wide, had recently moved upward to a position 1,219 meters (4,000 feet) below the 3,794-meter-high (12,448 feet) summit ([Zandomeneghi 2013](#)). The researchers' acoustic imaging of the volcano's internal structure revealed that it is a complex network of deep faults. These faults likely connect to and are part of the WARS.

The noticeable absence of ice and snow in large portions of the McMurdo Dry Valley (MDV) area (Figure 7) is attributed by the author to HCF. The authors' geological evaluation of MDV topography showed that the numerous long strongly linear valley trends and cross-cutting long strongly linear valley trends are the result of faulting and cross-faulting related to stresses associated with lateral movement of the WARS. Very high HCF in areas immediately adjacent to the MDV, most notably the Mount Erebus Volcanic Complex, are thought to be connected to and fueling the HCF which is melting the MDV snow and ice.

Recent research confirms that the Blood Falls feature in the MDV area is associated with distinct long linear bedrock trends ([Badgeley 2017](#)). These trends are interpreted to be faults associated with the WARS. These local Blood Falls faults are likely acting as conduits that tap into WARS heat flow and iron enriched heated fluid flow.

Figure 7 also illustrates numerous pieces of glacial ice floating in the bay area located between the Mount Erebus High Bedrock Heat Flow Area and the likely high heat flow Pluton Area. This includes floating ice from the Nansen Glacier. This specific glacier has been the focus of numerous research and media articles stating that breakup of this glacier is the result of anthropogenic warming. However, given that; high HCF from the Mount Erebus Volcanic Complex and the Pluton Area are present in the immediate bay area, the Nansen Glacier lies directly atop a major WARS fault segment, and that the Nansen Glacier Valley is a long linear feature it is far more likely that Nansen Glacier ice breakup is related to HCF from this entire region.



Figure 7.) Mount Erebus high bedrock heat flow area located along a very active segment of the West Antarctic Rift System. Note McMurdo Dry Valley Area (MDV) and volcanoes (red circles) (credit google Earth and J. Kamis).

[Kerguelen Volcanic Plateau](#)

Figure 3 illustrates the location of East Antarctica's offshore Kerguelen Volcanic Plateau. It is one of the largest igneous provinces in the world covering 1,243,190 square kilometers (480,000 square miles) and rising 2,011 meters (6,600 feet) above the surrounding ocean basins. This plateau has a long and complicated geological history involving multiple movements of major plate boundaries resulting in development of numerous major faults, terrestrial volcanoes, seamounts, hydrothermal vents, and multiple lava flow layers. Recent research has shown that numerous geological features associated with this vast ocean floor volcanic plateau are actively emitting significant amounts of HCF into surrounding ocean, land and atmosphere.

Australian Antarctic Research Institute studies concluded the following about the effect of geologically induced HCF into the oceans and onto terrestrial regions of the Kerguelen Plateau ([Australian Antarctic Division 2016](#) and [Kamis 2016](#)).

- The area is currently geologically active based on observation of a terrestrial volcanic eruption and discovery of numerous very active hydrothermal vents.
- The area's extensive reoccurring plankton blooms are strongly influenced by iron enrichment related to chemically charged heated fluid flow from the area's countless hydrothermal vents and seamounts.
- The alteration of marine animal migration patterns such as seals, whales, etc. were the result of volcanic activity and associated HCF into the area's ocean waters.

Multiple terrestrial and ocean floor volcanic eruptions since 1992 from the McDonald Islands area portion of the plateau, including the 2016 eruption of the area's Big Ben Volcano, indicate that significant portions of the Kerguelen Plateau are currently geologically active ([Smithsonian Institution GVP](#) and [Volcano Live](#)).

April 2016 research by Durham University in the far eastern portion of East Antarctica concluded that very long linear west to east oriented subglacial valleys are likely the result of faulting. These valleys form a subglacial drainage system of numerous interconnecting streams and one large lake all of which flow downhill into the adjacent ocean ([Jamieson 2016](#)). The author believes that the existence of a subglacial drainage system in this area is evidence of HCF and that the faults may be a westward extension of the Kerguelen Volcanic Plateau ([Kamis 2016](#)).

[Weddell Sea](#)

On or about September 9, 2017 a huge sea ice melt hole (polynya) formed in a very thick portion of Antarctica's Weddell Sea Ice (Figure 8). Scientists at the Helmholtz Centre for Ocean Research attributed this melt event to periodic unexplained upward movement of bottom hugging very warm seawater ([GEOMAR 2017](#)). Others have speculated that the polynya formation involved lateral movement of deep warm bottom hugging seawater. Once this laterally moving seawater intersected the Maud Rise bathymetric high it was deflected upward and onto the base of the sea ice thereby melting the overlying sea ice ([Swart 2017](#)).

This interpretation has numerous problems as follows:

- The Maud Rise is more correctly termed the Maude Volcanic Plateau, because it is the result of lava flow layering ([Schandl 1990](#)) and the complex interaction of ocean floor faulting and associated seamounts (J. Kamis observations).
- The other bathymetric highs in this portion of the Weddell Sea are not associated with sea ice melting.
- The sea ice melt hole (polynya) exactly matches the outline of the Maude Volcanic Plateau and does not elongate down current from the bathymetric Maud high (Figure 8).
- The warm sea area is associated with the Maud Volcanic Plateau (Muench 2001).
- Most importantly, the melt hole appeared in 1976 and then did not reappear until 2017 ([GEOMAR 2017](#)). This is characteristic of the erratic eruptive nature of volcanic heat flow pulses, and not characteristic of moderately rising atmospheric temperatures.



Figure 8.) Bathymetric and geological setting of the Maud Volcanic Plateau Region. Sea ice melt hole outlined in red. (credit Google Earth and J. E. Kamis).

An alternative and data supported explanation of the Weddell Sea polynya associated with the ocean floor Maud Volcanic Plateau is generation by HCF emanating from active geological features present on the plateau surface.

In summary, most of the bedrock geological features of West Antarctica have in past and still to this day actively flow significant amounts of HCF onto the base of the ice sheet and into adjacent oceans. These emissions are the dominant force acting to alter Antarctica's; ice sheet mass and extent, adjacent ocean water temperatures, sea ice composition and extent, plankton bloom vitality, and marine animal migration patterns. In stark contrast, most of East Antarctica's bedrock geological features are currently non-active. In very telling fashion, this portion of the continent is not experiencing the type of significant changes observed in West Antarctica. All this information demonstrates that geological and not manmade atmospheric forces are the dominant cause of observed changes in the Antarctic Continent in accordance with Plate Climatology Theory

Arctic Region

The Arctic region's glacial ice sheets, sea ice, surrounding ocean waters, biological regimes, and lower atmosphere are currently undergoing significant change. Scientists advocating anthropogenic warming have championed these changes as proof positive that anthropogenic warming is the root cause of the changes. Implicit in their argument is the notion that the Arctic's geological features have little or no influence on this region's climate and climate related events presumedly, because they envision these geological features as; inactive, slightly active, active within limited areas, or emitting minimal amounts of heated chemically charged fluid. The author presents data in this section that demonstrates that geological processes are an underappreciated contributor to the modern-day changes seen in the Arctic.




Figure 9. Arctic region active geological features (black cross-hatched) and referenced very high heat flow areas (red cross-hatched) (credit Google Earth and J. E. Kamis).

Significant amounts of data, numerous research studies, and observations by the author indicate that many of the Arctic's geological features are in fact very active and emitting massive amounts of HCF into Arctic region oceans, bedrock, and atmosphere ([Carmack 2012](#)). Evidence supporting this idea is abundant, reliable, and convincing.

The major geological features of the Arctic region as shown on Figure 9 are:

- Greenland / Iceland Mantle Plume
- Mid-Arctic Rift System
- Baffin Bay / Labrador Rift System
- Aleutian Island / Kamchatka Convergent Plate Boundary

All these geological features are proven emitters of significant amounts of HCF. Specific examples of emissions from these major geological features and their resulting climate effect are discussed below.

[Greenland - Iceland Mantle Plume](#)

Present day accelerated melting of the Greenland ice sheet and associated sea ice is primarily the result of two factors. Geologically induced warming of oceans adjacent Greenland, and most importantly relic heat flow from the Greenland / Iceland mantle Plume.

Figure 10 illustrates ancient position of and present-day relic heat flow the Greenland / Iceland Mantle Plume:

- North-northwest movement of Greenland across the stationary Greenland / Iceland Mantle Plume during the last 100 million years ([Martos 2018](#)). The heavy black dashed line illustrates the tract of this movement. Black dots mark the position and date of Greenland relative to the non-moving mantle plume.
- Greenland's current and anomalously high bedrock heat flow. Clearly relic heat from the mantle plume is still greatly affecting and elevating Greenland's current bedrock heat flow. The magnitude and extent of this relic heat flow has been confirmed by three other research studies ([Rezvanbehbahani 2017](#), [Rogozhina 2106](#), [Fahnestock 2001](#)). All researchers conclude that this relic heat flow is contributing significantly to basal melting of Greenland's Ice Sheet. It is also possible that outflow from the heated basal meltwaters are acting to warm ocean waters in proximity to the outflow points further exacerbating ice sheet melting.
- Rapid retreat of eastern Greenland's Young's Valley Fiord (Figure 10) is a specific example of how geologically induced bedrock heat flow directly affects the melting and lateral flow rate of individual Greenland glaciers.




Figure 10.) Ancient tract and present-day heat flow of Greenland / Iceland Mantle Plume (map credit Martos 2018, some of the labels J. E. Kamis).

Researchers from Aarhus University concluded that geothermal heat flow was accelerating the melting and lateral flow rate of the Young's Sound Glacial Valley (Figure 10 and quote below from [Rysgaard 2018](#)).

"Currently, there is a large disagreement between observed and simulated ice flow, which may arise from inaccurate parameterization of basal motion, subglacial hydrology or geothermal heat sources. Recently it was suggested that there may be a hidden heat source beneath GIS caused by a higher than expected geothermal heat flux (GHF) from the Earth's interior. Here we present the first direct measurements of GHF from beneath a deep fjord basin in Northeast Greenland. Temperature and salinity time series (2005–2015) in the deep stagnant basin water are used to quantify a GHF of $93 \pm 21 \text{ mW m}^{-2}$ which confirm previous indirect estimated values below GIS. A compilation of heat flux recordings from Greenland show the existence of geothermal heat sources beneath GIS and could explain high glacial ice speed areas such as the Northeast Greenland ice stream."

The greater Young's Valley Fjord region is home to long linear active hydrothermal vent complexes, ocean floor hot vents, valley trends, and offshore fault trends ([Rysgaard 2018](#) and J. E. Kamis observations). The author speculates that the heat source fueling the hydrothermal vent complexes and hot vents is being supplied by the Mid-Arctic Rift System. Heat of the Young's Valley Fjord bedrock is likely from two sources, the Mid-Arctic Rift Fault System and relic heat flow from the Greenland / Iceland Mantle Plume.

[Mid-Arctic Rift System](#)

Numerous geological features associated with the Mid-Arctic Rift System have been shown to be emitting significant amounts of HCF into Arctic ocean waters as per the following examples.

The Jan Mayen Volcanic Complex is a 1,500 kilometer (932 mile) long seafloor segment of the Mid-Arctic Rift System (Figure 9). This complex is home to numerous active; hydrothermal vents, seamounts, and faults. It is prone to numerous earthquakes testifying to its active geological nature as per a November 9, 2018 6.8 Richter scale earthquake located in close proximity to Jan Mayen Island. Researchers from the University of Bergen led by Dr Pedersen had this to say about the Jan Mayen geologic feature:

"2,200 F magma pouring into the seas from hundreds of submarine volcanoes – and we wonder why the seas are warming. We have found volcanoes at such a shallow level and they could break the surface at any time and form a new island group. Just for the record, I've been talking about underwater volcanoes for years. In fact, there's an entire chapter in my book "Not by Fire but by Ice" that discusses the importance of underwater volcanoes, and how they're heating the seas" ([Ice Age Now 2013](#), [Pedersen 2013](#), and [Olsen 2014](#))

The northern end of the Jan Mayen Volcanic complex includes and is defined by the Svalbard Islands. These islands are currently geologically active as indicated by; significant emissions of hydrothermal methane ([Mau 2107](#)), differential melting of its ice sheets ([Kamis 2015](#)), and numerous earthquakes ([University of Norway 2013](#), [Kvaerna](#), and [Antonovskaya 2015](#)). The long linear trends of Svalbard area seafloor methane emissions perfectly match the orientation of known seafloor faults that are part of the Mid-Arctic Rift System. This is strong indication that the origin of the methane emissions is hydrothermal, geologically induced heat flow melting of methane clathrate beds, or a combination of both. Melting of one Svalbard Island ice cap, Austfonna, suddenly accelerated in 2012. No other Svalbard Island ice caps, including the adjacent Vestfonna ice cap, experienced increased melting [Kamis 2015](#) and ([McMillan 2014](#)). This differential ice cap melting is very likely related to differential HCF from faults related to the Mid-Arctic Rift System which dissect the Svalbard Islands. Researchers have noticed that in recent decades that areas north of the Svalbard islands have been ice free for longer periods of time ([Polyakov 2017](#)). This differential sea ice melting is also likely related to HCF.

In April of 2014 a 4.5 scale earthquake occurred along a segment of the seafloor Mid-Arctic Rift System. This quake was likely triggered by magma chamber movement beneath the Mid-Arctic Rift. The shifting magma chambers released heat and hydrothermal methane into the deeper extent of rift faults. This heat and methane

moved upward along the faults and the process acted to melt additional shallow layers of frozen methane locked in methane clathrate beds adjacent to the faults. The upwardly migrating heat and methane was released into the overlying ocean acting to warm the water column. The methane continued to rise through the ocean column and into the atmosphere where it was recorded by NASA satellites ([Kamis 2015](#)). Worldwide there are multiple proven examples of massive methane pulses associated with seafloor earthquakes ([Fischer 2013](#) and [Hokugo 2018](#)).

On October 11, 2015 a melt hole rapidly developed in a very thick portion of the Arctic Sea Ice ([Kamis 2015](#)). The sea ice melt hole was positioned directly above the seafloor location of the currently active Jessica's Hill Vent Complex. The limited size, rapid occurrence, and position above a known hydrothermal vent complex of this melt hole is strong evidence that it is geological in origin.

Ultra-slow-moving rift systems such as the Mid-Arctic Rift System have been the focus of numerous research studies for many years, such as but not limited to, [Schlindwein 2005](#) and [Schlindwein 2016](#). Most of these studies have concluded that ultra-slow-moving rift systems circulate seawater downward along system faults to a depth of 15 kilometers (9.3 miles) or more. Where it is heated and chemically charged, then pulsed upward back into overlying ocean waters. The emission magnitude and effect of these giant ocean circulation systems is not currently well known. However, the author theorizes that these systems are significant contributors of heat and methane into overlying ocean waters thereby having a strong influence on the regions oceans, atmosphere, and climate.

[Baffin Bay/Labrador Fault System \(BB/L\)](#)

The April 2014 4.5 Mid-Arctic Rift System earthquake apparently had effect on the Baffin Bay Labrador Rift System (Figure 9) because it also released a large pulse of methane into the atmosphere immediately after the earthquake ([Kamis 2015](#)). The author believes that this seafloor methane release is the result of a HCF pulse from the Baffin Bay / Labrador Rift system.

[Nares Strait Fault System](#)

Recent research and observations by the author indicate that geological features associated with northwest Greenland's Nares Strait Fault System emit high amounts of HCF into the area's oceans, base of ice sheets, and atmosphere. Evidence supporting this idea is as follows.

Northwest Greenland's long linear Nares Strait ocean passageway is the surface expression of the 370-kilometer offset left lateral Nares Strait Fault System. This northeast to southwest trending fault system connects to the Baffin Bay spreading center fault system (Figure 9). The Nares Strait Fault System is home to several recently discovered high heat flow features.

First, two long linear and parallel Devon Island subglacial liquid water lakes. Devon Island is located at the structurally complex intersection of the Nares Strait and Baffin Bay fault systems. A recent research study discovered two long linear and very salty sub-glacial lakes on Devon Island. The orientation of the long linear subglacial lakes perfectly matches the trend of known surface faults associated with stresses from the left lateral Nares Strait Fault, including surface faults located at the edge of the glacial ice. Knowing that the overlying glacial ice is composed of freshwater, the high salt concentration of the subglacial lakes implies an HCF cause for lake development ([Rutishauser 2108](#) and [Kamis 2018](#)).

Second, significant melting and retreat of one of two large glaciers located in northwest Greenland that are a mere seven miles apart ([Kamis 2018](#) and [NASA](#)). Recent NASA research showed that the Tracy Glacier has been bottom melting and retreating to the east, while the adjacent Helprin Glacier has not retreated. The NASA study also showed that the bay water beneath the Tracy Glacier is anomalously warm, while bay water beneath

the Helprin Glacier is of normal temperature which is a common occurrence in geothermal heat sourcing configurations involving complex fault systems. The orientation and long linear shape of both the Tracy and Helprin Glacial valleys is strong evidence that they are fault induced features owing their generation to stresses associated with the left lateral Nares Strait Fault System.

Lastly, the rapid melting of the ice covering the world's largest high arctic lake, Lake Hazen [Lehnherr 2018](#)). Lake Hazen is located on Ellesmere Island which forms the northern western boundary of the Nares Strait ocean passageway. A passage way which was formed by movement of the Nares Strait / Wegener left lateral fault system (Pipejohn 2008). Lake Hazen is located within this tectonically active region. The lake owes its very existence to fault movement of the Lake Hazen Thrust which acted to down drop and form an enclosed long linear valley. The Lake Hazen thrust connects to Nares Strait / Wegener Left Lateral Fault System at an angle that implies it is part of this fault system (Pipejohn 2008). Indications of high HCF in the Lake Hazen area include:

- Structural association with the Nares Strait / Wegener Left Lateral Fault System that is thought to be a source of HCF ([Kamis 2018](#)).
- Proximal position to mantle plume relic heat flow and a position directly atop the ancient path of the Greenland / Iceland Mantle Plume.
- Proximity to mineral rich springs. There are a total of four mineral-rich springs located in the Lake Hazen region ([Sapers 2017](#), [Pollard 1998](#), and [Omelon 2006](#)). The author believes these springs are associated with faults which tap into relic mantle plume heat flow. Although the temperature of the spring's surface water is cool, it is likely that at slightly deeper depths along the sourcing fault planes the spring's water is warmer.
- The unusually hot spring-like mineral and biological content of these spring's surface water ([Kamis 2018](#))

[Aleutian Island Convergent Fault System](#)

The 4,000-kilometer-long (2,485 mile) Aleutian Island Convergent Fault System forms the southern and enclosing boundary of the Bering Sea (Figure 9). A review of historical Sea Surface Temperature Maps of the Bering Sea show that starting on or about 2012 and continuing through the present-day seawater temperatures in this area are greatly elevated.

The Aleutian Island Convergent Fault System is home to an estimated 80 active or semi-active terrestrial volcanoes. Based on bedrock heat flow information, the Aleutian Island Convergent Fault System seafloor regions likely contain a very large number of seamounts, active faults, and hydrothermal vents ([Batir 2013](#) and J. Kamis observations). These geological features and others scattered across the floor of the Bering Sea are likely emitting large amounts of HCF into Bering Sea and warming this ocean region as per analysis of NOAA Shallow Sea Surface Temperature Maps (J. Kamis observations). Once warmed ocean currents carry this warmed seawater northward eventually funneling it through the Bering Strait and into the western Arctic Ocean where it acts to warm seawater in this area and melt the western edge of the Arctic Sea Ice ([Kamis 16 July 2018](#)). Recent research by Yale and Woods Hole Oceanographic Institute is interpreted to support a Bering Sea heating source for a just discovered very large bottom hugging warm pool located in the western Arctic Ocean ([Timmermans 2018](#)).

Evidence that additional warming of western Arctic Ocean waters is from active seafloor geological features comes from a study by Scripps Institution of Oceanography. The study shows that massive amounts of turbulent upwelling very warm ocean water likely originates from the Arctic seafloor ([MacKinnon 2015](#)). Lead scientist Dr. Jennifer MacKinnon said this:

"The strength of heat coming up from below the surface has been as strong as the heat coming down from the Sun,...There's a reservoir of heat in the Arctic Ocean, well beneath the surface,

that historically – when there's been a lot of ice has been fairly quiescent,....The heat is being brought to the surface by surprisingly strong eddies,..."The strength of [these currents] has been incredible."

Dr McKinnon and her research team theorize that strong surface winds are acting to generate large turbulent ocean cells that bring very warm bottom hugging ocean water to the surface where it acts to melt Arctic sea ice. The author theorizes that geologically induced seafloor heat flow from active geological features is generating the deep ocean warm pool and creating the turbulent surface reaching eddies.

In summary, the Arctic region is home to numerous active geological features that flow significant amounts of HCF into adjacent oceans and the base of glacial ice sheets. These emissions are the dominant force acting to alter ice sheet mass and extent, ocean water temperatures and chemistry, sea ice thickness and extent, and marine animal migration patterns. All this information demonstrates, as per Plate Climatology theory, that geological and not manmade atmospheric forces are the root cause of observed changes in the Arctic region.

Oceans

Geologically induced HCF from seafloor and terrestrial geological features have in past and still to this day act to significantly alter many aspects of earth's oceans. Altered ocean water then acted, and still to this day acts to significantly influence earth's climate and climate related events. The author believes that due consideration should be given to geologically induced HCF as a possible explanation of present-day; slight decreases in ocean PH, some of the localized oxygen concentration decreases, and some of the decreases in marine animal populations. Evidence supporting these statements is as follows.

Present day changes of PH in several of earth's oceans has been termed "Acidification". However, most research indicates that during the last 300 million years none of earth's oceans have ever had, nor do they currently have a PH equal to or less than 7. Therefore, earth's oceans are not, nor have they for millions of years been acidic. To the contrary, they have varied from basic to slightly less basic, ranging from an estimated ancient PH of 8.2 to a current PH of approximately 8.1 ([Nat Geo 2017](#) and [Hönisch 2012](#)).

A specific example of ancient ocean PH alteration by geological forces comes from research of the Paleocene-Eocene Thermal Maximum event. This event occurred 56 million years ago and was the result of a huge worldwide volcanic pulse. A volcanic pulse which acted to infuse the atmosphere with a much greater concentration of CO₂, lower the PH of earth's oceans, and significantly alter ocean biologic environments ([Keller 2018](#)).

Another example of how geological forces can alter earth's oceans are ancient worldwide anoxic events that acted to dramatically reduce ocean oxygen content leading to mass extinctions. Several research studies have concluded that the Toarcian Oceanic Anoxic Event (TAE), which occurred 183 million years ago, was generated by a huge pulse of volcanic activity that acted to infuse earth's atmosphere with massive amounts of CO₂. This atmospheric CO₂ then acted to de-oxygenate earth's oceans and force a worldwide mass extinction ([Them 2018](#) and [Zheng 2013](#)). The Ocean Anoxic Event (OAE) occurred 94 million years ago and was fueled by a large volcanic emission pulse ([Clarkson 2018](#)). Recent research has shown that the major cause of the KT extinction event was volcanism within India's Deccan Traps area ([Keller 2018](#)). An event that effected both oceans and atmosphere.

Pacific Ocean

[El Nino](#)




Figure 11. 1997 El Nino Pacific Ocean warming event. Red shades indicate anomalously warm ocean water. (image credit NASA/Goddard Space Flight Center and label credit J. E. Kamis)

El Nino is one of the most influential climate and climate related phenomena on earth. These massive pulses of warm seawater have in past and still to this day act to dramatically alter many aspects of earth's oceans (temperatures, chemistry, currents, and sea level) and atmosphere (temperatures, chemistry, and winds). These alterations significantly influence; ocean biological regimes, terrestrial biological regimes, human activities, local weather, and worldwide climate. Some attribute the sudden migration of ancient human civilizations to El Nino induced climate changes.

The generation and maintenance of El Nino's involves the complex and unique interaction of geological forces with deep ocean waters in a geographically limited seafloor location of the far western Pacific Ocean (Figure 12, [Kamis 2014](#), [Kamis 2015](#), and [Kamis 2017](#)).

El Nino generation is preceded by pulses of earthquake activity associated with major deep reaching faults, many of which are plate bounding faults, in a geographically limited location (Figure 12). This tectonic activity acts to alter the permeability and hydrologic fluid flow characteristics of deep ocean floor rock layers adjacent to the tectonically active ocean floor faults. Fault movements that fracture ocean floor brittle rock layers for a great distance away from the fault plane, does not fracture interbedded ductile rock layers rock layers, and connects the now high permeability fault plane to underlying magma chambers. The result of this activity is to

generate a large and very powerful seafloor mega-fluid flow system ([Baker 1987](#)) that emits massive amounts of HCF into overlying ocean waters ([Kamis 2015](#)).

Movement of fluid through the now activated El Nino ocean floor mega-fluid flow system is driven by downward pressure from the overlying ocean column. This pressure forces seawater into the open fault plane eventually placing it in contact with underlying magma chambers where it is heated and chemically charged. The altered seawater is then forced laterally into and constrained within fracture permeable rock layers. As the altered seawater moves laterally farther and farther away from the fault plane it eventually encounters vertical permeability pathways allowing it to flow upward and back into the ocean where it acts to heat and chemically charge ocean water thereby generating and maintaining El Nino's.

A summary of evidence that supports El Nino generation by geological induced heat flow and chemically charged heated fluid flow is as follows:

- All El Niño's have originated at the exact same deep ocean seafloor geographically limited fixed area located east of the Papua New Guinea / Solomon Island area. This is one of the most volcanically and seismically active areas on earth. The conjunction of five major plate boundaries, source point for earth's second largest seafloor volcanic plateau, thousands of seamounts, countless numbers hydrothermal vents, and a complex network of significant faults (Figure 12).
- El Niño sea surface temperature and chemical anomalies have linear and intense boundaries inferring that the energy source is stationary and very powerful (Figure 11 and [Kubota 2014](#)).
- The shape of all El Niño sea surface temperature anomalies are identical, and not like other sea surface temperature anomalies.
- The fixed source point and cone shape of heat and ash emissions from land based volcanic eruptions are an excellent analog to the cone shape of heat emissions all El Nino's. Additionally, the cone shape of deep-ocean geological hydrothermal vent emissions is another very good mini-analogy of the larger El Niño's.
- El Niño's occurrences do not correlate with atmospheric changes.
- El Nino's often occur in "bundles". Multiple El Nino's occur in a short time span, grouped together. Typically, the first El Nino in a bundle is of lower intensity, subsequent El Nino's are progressively more intense, often ending with a final maximum/high intensity El Nino. This El Nino bundle pattern is very similar to the progression of well monitored and well understood land based volcanic and tectonic events which typically build through time to a final large volcanic eruption or tectonic event.
- El Niño-like events do not occur elsewhere in Pacific. Why? If they are atmospheric in origin there should at least be another mini-El Niño's. There are none.
- Historical records indicate that the first "recorded" El Niño occurred in 1525 as observed by Spanish explorers. Other studies suggest strong ancient El Niño's ended the Moche Peruvian Civilization ([Etayo-Cadavid 2013](#)). The main point here is that strong El Niño's are natural, and that they are not increasing in relationship to anthropogenic warming.
- El Nino ocean warming acts to strongly effect ocean coral reef systems, often referred to as "Coral Bleaching". The author believes that these alterations are natural and necessary effects of geological forces. Natural effects which fit into Natural Selection Theory developed by Geologist Charles Darwin in the geologically active Galapagos Island Rift System region ([Kamis 2015](#) and [Kamis 2016](#)).
- El Nino warming and chemical changes of Pacific Ocean seawater has a strong influence on Pacific Ocean phytoplankton distribution ([NASA 2009](#) and
- La Nina's cool ocean water flows originate at the same fixed-point source as El Niño's indicating fluid emissions from a fixed geological feature.
- El Niño / La Nina prediction models loaded with atmospheric and shallow oceanic data consistently fail, likely because they are modelling the "side effects" of geologically warmed / cooled oceans and not the "cause" of the El Niño / La Nina event.

- El Nino / La Nina events are associated with seismicity and, or volcanism in the point source area ([Kamis 2017](#), and [Guillas 2010](#)).

La Nina

The generation and maintenance of La Nina's also involves the complex and unique interaction of deep ocean and geological forces in one specific geographically limited seafloor location in the far western Pacific Ocean. Specifics of exactly how geological forces generate and maintain La Nina cooling is not well understood. The author believes that el Nino phase of the El Nino / La Nina geological phenomenon develops as follows.

Once the heating power of the still actively flowing El Nino Mega-Fluid Fluid Flow System wains it morphs into a cooling phase. This is due to the progressive conversion of system free-flowing hydrothermal methane to frozen methane clathrate along the fault plane and within adjacent permeable rocks layers. A conversion process that acts to quickly cool system seawater and overlying ocean waters. Eventually frozen methane clathrate emplacement completely fills all permeability within fault planes and fractured rock layers thereby ending the La Nina phase and the El Nino / La Nina cycle.

El Nino / La Nina Seafloor Source Point

Scientists have failed to locate the fixed seafloor geological heat and cooling source point of the El Niño / La Nina events for two reasons, insufficient data and “Atmospheric Bias”.




Figure 12.) Bathymetric map of the western Pacific Ocean with El Nino Heat Source Area (Bathymetric map credit Schmidt Ocean Institute, credit "El Nino Heat Source Area" outline and label J. E. Kamis).

Concerning insufficient data, the earth's oceans cover 361,821,339 square kilometers (139,700,000 square miles) of our planet's surface, however as per NOAA there are only 3,908 floats in the worldwide Argo Buoy System that record shallow to upper mid-level ocean temperatures. This equates to one buoy every 117,000

square kilometers (45,174 square miles). Furthermore, the oceans can be generally divided into three depth range layers; shallow, mid-level and deep. So, at any time the buoys may be stationed at different ocean levels. Therefore, the weekly or monthly amount of buoy measurements per square mile within a specific ocean layer may be less dense than the 117,000 per square kilometer (45,174 square mile) average. Many deep-ocean geological heat and fluid sources are located at depths greater than 2,000 meters (6,562 feet) which is beneath the ARGO Buoy depth limits. The Pacific Ocean has an average depth of 4,270 meters (14,000 feet) and the El Nino / La Nina source point area (Figure 12) is at a depth of 3,660 meters (12,000 feet). Therefore, the buoys are incapable of locating or precisely measuring heat flow from deep Pacific Ocean geological features. The El Nino heat source area represents 233,000 square kilometers (86,100 square miles) which is a mere 0.01 percent of the Pacific Ocean's 166,000,000 square kilometer (64,092,958 square mile) area (Figure 12). The Argo Buoy System could easily miss this 0.14 percent 3,658 meter (12,000 feet) deep geological heat source point area.

Atmospheric Bias has played a large role in preventing scientist from discovering the origin of and locating the heat source point of El Nino warming. Lacking large amounts of HCF emission data from seafloor geological features in the western Pacific Ocean scientists looked to the immense atmospheric and shallow ocean emission data set for answers concerning what force or forces generate and maintain El Nino's. Scientists did in fact find correlations between atmospheric and shallow ocean data with many aspects of El Nino generation and maintenance. However, the author contends that these correlations are not cause and effect, rather cause and side effect. Once geological forces act to alter ocean temperature, currents, and chemistry, the changed ocean then acts to alter properties of the overlying atmosphere such as trade winds, storm activity, temperature patterns, and precipitation. These oceanic and atmospheric changes are side effects of geological forces. This bias towards atmospheric and shallow ocean data is here termed "atmospheric Bias".

Arctic Ocean

Arctic Amplification

Arctic Amplification which is generally defined as changes in net radiation balance such as intensification of greenhouse gas related warming, has been a long-standing tenet of the Global Warming / Climate Change Theory. However new research ([Stuecker 2018](#)) concludes that many aspects of this tenet are incorrect or in need of significant modification as follows:

- Increased atmospheric CO2 concentrations and resulting Arctic atmospheric warming are not the result of worldwide CO2 sources, rather the result of local CO2 sources.
- Warming of the Arctic's lower atmosphere does not mix with the overlying or lateral atmosphere, rather is strongly restricted vertically and geographically.
- The "Ice Albedo Affect" does not significantly contribute to Arctic atmospheric warming, rather has an insignificant warming effect.
- Worldwide cloud, ocean current, or hot winds such as those related to El Nino do not significantly contribute to Arctic region atmospheric warming, rather have little or no effect.

These conclusions indicate that Arctic Amplification is the result of local, not worldwide forces. The author believes that these local forces and resulting changes to the Arctic atmosphere are the result of seafloor HCF from two active Arctic geological features, the Aleutian Island Subduction Zone and the Mid-Arctic Rift System.

As previously discussed in this paper, these two active geological features are thought to be emitting significant amounts of HCF into the Arctic Ocean thereby warming, in varying degree, the entire Arctic seawater column. This warmed seawater then acts to significantly warm the overlying Arctic atmosphere especially in areas of the atmosphere located in close geographical association with HCF ocean warm pockets.

Figure 13, taken from NOAA's 2018 Arctic Report Card, clearly shows that:

- The extent of recently elevated Arctic atmospheric temperatures closely matches the outline of the Arctic Ocean.
- The configuration of the western intensely atmospherically warmed area can be interpreted to be the result of HCF warmed seawater inflow from the Bering Sea into the Chukchi Sea.
- The configuration of the eastern atmospherically warmed area can be attributed to HCF from the underlying Mid-Arctic Rift System.

The above information supports the idea that Arctic Amplification is largely the result of geological and not atmospheric forces.




Figure 13.) Arctic region difference from average lower atmosphere temperature October 2017–September 2018 (map credit NOAA, HCF lines and labels J. Kamis).

North Atlantic Ocean

Gulf Stream "2010 Slowdown"

The strong climate influencing Gulf Stream Current slowed down in May 2010. This significant ocean current event was given many names, "Gulfstream Shutdown", "2010 Slowdown", etc. Most climate scientists attributed the root cause of this event to the general effects of anthropogenic warming. Others attributed this slowdown to the effects of a stationary shallow water North Atlantic warm water cell of unspecified origin that acted to block or inhibit eastern flow of the Gulf Stream current ([Buckley 2015](#)).

The author believes that this blocking warm cell was generated by a massive pulse of HCF from ocean floor rift faults ([Kamis 2015](#)). There are several lines of evidence that support this claim. The outline and geographic position of several major North Atlantic seafloor faults perfectly matches the outline and geographic position of

the overlying shallow water warm cell. This is compelling evidence of a cause and effect relationship between fault heat flow and warm cell generation. It is thought that a massive long-lasting pulse of heat from seafloor faults rose vertically through the ocean column and acted to warm shallow ocean water directly above the faults thereby forming the blocking warm cell. Additionally, the warm cell has a distinct and well-formed outline that developed rapidly. This is very characteristic of the erratic and sudden development of geological features. Lastly, the stationary position of the warm cell implies a powerful and non-moving heating mechanism, again characteristic of a geological feature.

Terrestrial Volcanic Regions

Quantifying the climate effect of terrestrial volcanic emissions is a complex and difficult task. Most researchers have focused their attention on detailing the climate effects of major world class volcanic eruptions. The climate effect of these events on ocean, atmosphere, and climate have been well documented, especially during the last 5-10 years. However, several aspects of the climate effect of terrestrial volcanic eruptions has been lost in the shuffle, specifically the:

- Climate effect of CO₂ and other gas emissions from non-erupting but active volcanoes.
- Climate effect of CO₂ and other gas emissions from minor volcanic eruptions.
- Science behind distinguishing the difference between volcanic emissions and emissions from burning of fossil fuels.

Non-Erupting Volcanic CO₂ Emissions

Just released research of Iceland's non-erupting Katla Volcano indicates that scientists have greatly underestimated the emission amount and significance of non-erupting terrestrial volcanoes ([Ilyinskayasee 2018](#)). The study concluded that Katla is emitting massive amounts of CO₂ into the atmosphere.

"The emissions are in the order of 12 to 24 kilotons a day which means 4 to 8 megatons a year. That's the same output as an extra 1 – 2 million cars on the road each year. Ilyinskaya et al, Geophysical Research Letters We discovered that Katla volcano in Iceland is a globally important source of atmospheric carbon dioxide (CO₂) in spite of being previously assumed to be a minor gas emitter. Volcanoes are a key natural source of atmospheric CO₂". ([Ilyinskayasee 2018](#))

Minor Eruptive Volcanic CO₂ Emissions

Results of a 2014 research study by MIT concerning the climate influence of volcanic eruptions concluded that volcanic emissions greatly effect climate and climate related events. MIT said that the global warming hiatus was forced in part by the cooling effect of volcanic eruptions and that most climate models have not accurately accounted for the effects of volcanic activity ([MIT 2014](#)). Other MIT research entitled " *Small Volcanic Eruptions could be Slowing Global Warming*" ([MIT 2014](#)) speaks to the underappreciated climate effect of eruption emissions from earth's numerous small volcanic features.

Volcanic vs Fossil Fuel CO₂ Emissions

Some have presented convincing evidence that natural volcanic and man-made CO₂ emissions have the exact same and very distinctive carbon-14 isotopic fingerprint ([Casey 2014](#)). Scientists advocating the GWCC theory believe, based on supposedly reliable research (Gerlach 2011), that volcanic emissions are minuscule in comparison to human-induced CO₂ emissions. As a result, they assert that modern day volcanic emissions have little or no effect on climate. I believe that these assertions are based on less than sufficient evidence. Gerlach's volcanic CO₂ calculation was based on just 7 actively erupting land volcanoes and three actively erupting ocean floor hydrothermal vents (seafloor hot geysers). Utilizing gas emission data from this very limited number of volcanic features, Gerlach estimated that the volume of natural volcanic CO₂ emissions is 100 to 150 times less than the volume of man-made CO₂ emissions from the burning of fossil fuels and therefore of no consequence. To put this calculation process into perspective, the Earth is home to an

estimated 1,500 land volcanoes and 900,000 seafloor volcanoes/hydrothermal vents. By sampling just an extremely small percent of these volcanic features the data is insufficient to validate this calculation. Especially knowing that volcanic activity varies greatly from area to area, volcano to volcano, and through time. Utilizing just 0.001 percent of Earth's volcanic features to calculate volcanic CO2 emissions does not inspire confidence in the resulting value.

So-Called Ice Ages




Figure 14.) Major deep inner Earth reaching breaks in Earth's outer crust (spreading centers) are shown as black lines surrounded by red and yellow shades. Shading indicates the age of the ocean floor lava flows that are expelled from the spreading centers. Estimated maximum extent of glacial ice during glacial periods is shaded white and cross-hatched light blue (credit Estrada et. al. 2013, labeling and Ice Sheets by J. Kamis).

The very rapid and nearly instantaneous end of so-called [Glacial Periods](#) is caused by massive, short-lived pulses of super-heated seawater emitted in large part from major ocean faults termed [Tectonic Spreading Centers](#). Emissions that act to very rapidly warm and chemically alter oceans and atmosphere thereby melting glacial ice sheets.

For many years now accepted scientific theory has stated that there are distinct and separate Glacial and Interglacial Periods which when coupled together define one Ice Age. Solar and atmospheric forces are often cited as the root cause of these glacial and interglacial climate shifts.

Reinterpretation and then integration of information from the authors observations with two recent research studies indicates that this consensus theory is no longer correct.

Stated in another way, there are no such things as glacial periods, interglacial periods, or Ice Ages. Instead, just an extremely short time frame (1,000 – 3,000 year) major geologically induced worldwide heating event followed by a slower recovery to a here termed **Normal Climate Status**. During this **Recovery Period** earth's oceanic, terrestrial, and atmospheric regimes complexly interact until a Normal Climate Status is achieved typically within 75,000 years. Normal Climate Status is best described as; significant ice sheet extent across many continents, cooler ocean and atmospheric temperatures, and distinctly different biological regimes. Reoccurrence of major geologically induced heating events is associated with and generated by [Milankovitch Cycles](#) which occur approximately every 100,000 years. For a more detailed discussion of this new interpretation of so-called Ice Ages the reader is directed to [here](#), [here](#), and [here](#).

SUMMARY



Figure 14.) Schematic Illustration of Upper Mantle Convection Cell (Image credit public domain and label credit J. Kamis).

The present day and ancient climate effect of geological forces has been greatly underestimated and underappreciated. A more realistic assessment of their significant climate effect has been realized by integrating and then evaluating numerous examples of how geological forces effect, or have affected earth's oceans, atmosphere, biological regimes, and terrestrial areas. This assessment clearly shows that geological forces have a very significant, and in some cases, dominating effect on earth's climate and climate related events. Effects that have remained hidden for several reasons:

- Most of earth's active major geological features are in remote and difficult to research deep ocean or sub-ice sheet polar regions. As a result, their activity, function, and HCF are not well understood.
- Underestimation of CO₂ and methane emissions from terrestrial non-eruptive volcanic features and medium to small size mildly eruptive volcanic features.
- Incorrectly attributing many climate and climate related events to atmospheric or oceanic side effects when the root cause is related to geological forces.

- Unappreciation of the analog nature of past geologically induced extinctions, ocean anoxic events, and alterations of Polar Ice Caps.

Finally, The Plate Climatology Theory provides plausible alternative explanations for many our planets climate phenomenon, trends, and climate related events. It is deserves strong consideration by mainstream climate scientists.

*James Edward Kamis is a retired professional Geologist with 42 years of experience, a B.S. in Geology from Northern Illinois University (1973), an M.S. in geology from Idaho State University (1977), and a longtime member of AAPG who has always been fascinated by the connection between Geology and Climate. More than 14 years of research/observation have convinced him that the Earth's Heat Flow Engine, which drives the outer crustal plates, is an important driver of the Earth's climate as per his **Plate Climatology Theory**.*

REFERENCES

Nerilie Abrams, Robert Mulvaney, Francoise Vimeux, Steven Phipps, John turner, and Mathew England, Ocean winds keep Antarctica cold, Australia dry, 11 May 2014, Nature Climate Change 4, 564-569(2014), <https://www.nature.com/articles/nclimate2235> and <https://phys.org/news/2014-05-ocean-antarctica-cold-australia.html#jCp>,

Andrew Allibone, Simon cox, Ian Graham, Robert Smellie, Roy Johnstone, Simon Ellery, and Ken Palmer, Granitoids of the Dry Valleys area, southern Victoria Land, Antarctica: Plutons, field relationships, and isotopic dating, New Zealand Journal of Geology and Geophysics, 36:3, 281-297, DOI: 10.1080/00288306.1993.9514576, 23 March 2010, <https://www.tandfonline.com/doi/pdf/10.1080/00288306.1993.9514576>

Ando T, Yamamoto M, Tomiyasu T, Tsuji M, and Akiba S., Mercury distribution in seawater of Kagoshima Bay near the active Volcano, Mt. Sakurajima in Japan, , Bull Environ Contam Toxicol. 2010 Apr;84(4):477-81. doi: 10.1007/s00128-010-9946-7. Epub 2010 Feb 25. <https://www.ncbi.nlm.nih.gov/pubmed/20182698>

Dermot Antoniades, Santiago Giralto, Adelina Geyer, Antonio Alvarez-Valero, Sergi Pla-Rabes, Ignacio Granados, Emma Liu, Manuel Toro, John Smellie, and Marc Oliva, The timing and widespread effects of the largest Holocene volcanic eruption in Antarctica, *Scientific Reports* volume 8, Article number: 17279 (2018), https://www.nature.com/articles/s41598-018-35460-x?error=cookies_not_supported&code=dbd45781-1884-4442-9870-a7117dab1c39

Galina Antonovskaya, Yana Konechnaya, Elena Kremenetskaya, Vladimir Asming, Tormod Kvaerna, Johannes Schweitzer, and Frode Ringdal, Enhanced Earthquake Monitoring in the European Arctic, *Polar Science*, Volume 9, Issue 1, March 2015, Pages 158-167, <https://www.sciencedirect.com/science/article/pii/S1873965214000644>

Army Corps of Engineers, Aleutian Arc, <https://www.iwr.usace.army.mil/Missions/Coasts/Tales-of-the-Coast/Americas-Coasts/Alaska-Coast/Aleutian-Arc/>

Australian Antarctic Division, Volcanic hotspot may fortify ocean life, Australian Antarctic Magazine, issue 30, June 2016, <http://www.antarctica.gov.au/magazine/2016-2020/issue-30-june-2016/science/volcanic-hotspot-may-fortify-ocean-life>

Jessica Badgeley, Erin Pettit, Christina Carr, Slawek Tulaczyk, Jill Mikucki, Berry Lyons and Midge Science Team, An englacial hydrologic system of brine within a cold glacier: Blood Falls, McMurdo Dry Valleys, Antarctica, *Journal of Glaciology*, Volume 63, Issue 239, June 2017 , pp. 387-400, https://pdfs.semanticscholar.org/e882/4c1ba0c8b476fecff6d97fa8937420829eef.pdf?_ga=2.147764658.443811328.1546966450-1392082564.1546966450

Edward Baker, Gary Massoth and Richard Feely, Cataclysmic hydrothermal venting on the Juan de Fuca Ridge, *Nature* volume 329, pages 149–151, 10 September 1987, <https://www.nature.com/articles/329149a0#article-info>

P. F. Barker and I. A. Hill, Back-Arc Extension in the Scotia Sea [and Discussion], *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences*, Vol. 300, No. 1454, Extensional Tectonics Associated with Convergent Plate Boundaries (Mar. 26, 1981), pp. 249-262, <http://adsabs.harvard.edu/abs/1981RSPTA.300..249B>

D. H. N. Barker and J. A. Austin Jr., Crustal Diapirism in Bransfield Strait, West Antarctica: Evidence for distributed extension in marginal-basin formation, *Geology* (1994) 22 (7): 657-660., 1 July 1994 [https://doi.org/10.1130/0091-7613\(1994\)022<0657:CDIBSW>2.3.CO;2](https://doi.org/10.1130/0091-7613(1994)022<0657:CDIBSW>2.3.CO;2)

Valentina Barletta, Michael Bevis, Benjamin Smith, Terry Wilson, Abel Brown, Andrea Bordoni, Michael Willis, Shfaqat Abbas Khan, Marc Rovira-Navarro, Ian Dalziel, Robert Smalley, Eric Kendrick, Stephanie Konfal, Dana Caccamise, Richard Aster, Andy Nyblade, and Douglas Wiens, Observed rapid bedrock uplift in Amundsen Sea Embayment promotes ice-sheet stability, British Antarctic Survey, New Antarctic Heat Map Reveals Sub-Ice Hotspots, 13 November 2017, <http://science.sciencemag.org/content/360/6395/1335>

Joseph Batir, David Blackwell and Maria Richards, Final Report to the Energy Authority and Alaska Center for Energy and Power, SMU Geothermal Laboratory, 15 June 2013, http://acep.uaf.edu/media/57321/HFMAK_final-report.pdf

Heinz Blatter, Why an ice age occurs every 100,000 years: Climate and feedback effects explained, *Science Daily*, 7 August 2013, <https://www.sciencedaily.com/releases/2013/08/130807134127.htm>

Martha Buckley and John Marshall, Observations, inferences, and mechanisms of the Atlantic Meridional Overturning Circulation, *Reviews of Geophysics*, 17 December 2015, <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2015RG000493>

Christo Buizert, Vasileios Gkinis, Jeffery Severinghaus, Feng He, Benoit Lecavalier, Phillip Kindler, Markus Lauenberger, Anders Carlson, Bo Masson-Delmonte, James White, Zhengyu, Bette Otto-Bliesner, and Edward Brook, Greenland temperature response to climate forcing during the last deglaciation, *Science* 05 Sep 2014: Vol. 345, Issue 6201, pp. 1177-1180, DOI: 10.1126/science.1254961, <http://science.sciencemag.org/content/345/6201/1177>

Eddy Carmack, William Williams, Sarah Zimmermann and Fiona McLaughlin, The Arctic Ocean warms from below, *Geophysical Research Letters*, VOL. 39, L07604, doi:10.1029/2012GL050890, 7 April 2012, <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2012GL050890>

Mathew Clarkson, Claudine Stirling, Hugh Jenkyns, Alexander Dickson, Don Porcelli, Christopher Moy, Phillip Pogge von Strandmann, Ilsa Cooke, and Timothy Lenton, Uranium isotope evidence for two episodes of

deoxygenation during Oceanic Anoxic Event 2, PNAS, 5 March 2018, 115 (12) 2918-2923, <https://doi.org/10.1073/pnas.1715278115>.

Hugh Corr, First evidence of under-ice volcanic eruption in Antarctica, British Antarctic Survey News Release, 20 January 2008, <https://www.bas.ac.uk/media-post/first-evidence-of-under-ice-volcanic-eruption-in-antarctica/>

Maximilian De Vries, Robert Bingham, and Andrew Hein, A new volcanic province: an inventory of subglacial volcanoes in West Antarctica, University of Edinburgh, 2018, https://www.geos.ed.ac.uk/homes/rbingha2/48_2017_Vries.pdf

Eugene Domack, Scientists Discover Undersea Volcano Off Antarctica, National Science Foundation, 31 May 2004, <https://www.sciencedaily.com/releases/2004/05/040527235943.htm>

R. Dziadek, Karsten Gohl, and Norbert Kaul, Elevated geothermal surface heat flow in the Amundsen Sea Embayment, West Antarctica, Earth and Planetary Science Letters, Volume 506, 15 January 2018, Pages 530-539, <https://doi.org/10.1016/j.epsl.2018.11.003>

Michael Etayo-Cadavid, C. Andrus, K. Jones, and G. Hodgins, Marine radiocarbon reservoir age variation in *Donax obesulus* shells from northern Peru: Late Holocene evidence for extended El Niño, *Geology*, 41(5):599-602, April 2013, DOI: 10.1130/G34065.1

https://www.researchgate.net/publication/274438811_Marine_radiocarbon_reservoir_age_variation_in_Donax_obesulus_shells_from_northern_Peru_Late_Holocene_evidence_for_extended_El_Niño

Mark Fahnestock, Waleed Abdallati, Ian Joughin, John Brozena, and Prasad Gogineni, High Geothermal Heat Flow, Basal Melt, and the Origin of Rapid Ice Flow in Central Greenland, *Science* 14 Dec 2001: Vol. 294, Issue 5550, pp. 2338-2342, DOI: 10.1126/science.1065370, <http://science.sciencemag.org/content/294/5550/2338.full>

David Fischer, Jose Mogollon, Micael Strasser, Thomas Pape, Gerhard Bohmann, Noemi Fekete, Volkhard Spiess and Sabine Kasten, Subduction zone earthquake as potential trigger of submarine hydrocarbon seepage, *Nature Geoscience* volume 6, pages 647–651 (2013), 28 July 2013, <https://www.nature.com/articles/ngeo1886>

Andrew Fisher, Kenneth Mankoff, Slawek Tulaczyk, W Tyler, Neil Foley, and WISSARD Science Team, High geothermal heat flux measured below the West Antarctic Ice Sheet, *Science Advances*, 10 Jul 2015: Vol. 1, no. 6, e1500093, DOI: 10.1126/sciadv.1500093

Helen Fricker, Ted Scambos, Robert Bindschadler, and Laurie Padman, An Active Subglacial Water System in West Antarctica Mapped from Space, *Science* 16 Mar 2007: Vol. 315, Issue 5818, pp. 1544-1548 DOI: 10.1126/science.1136897

GEOMAR, Antarctica: Return of the Weddell polynya supports Kiel climate model, Science News, Helmholtz Centre for Ocean Research Kiel, 29 September 2017, <https://www.sciencedaily.com/releases/2017/09/170929093337.htm>

Terry Gerlach, Volcanic versus Anthropogenic Carbon Dioxide, *Eos*, Transactions, American Geophysical Union Volume 92, Number 24, 14 June 2011, <https://doi.org/10.1029/eost2011EO24>

Ben Gurion, New Antarctic rift data has implications for volcanic evolution, *Physics Org News*, August 21, 2018, <https://phys.org/news/2018-08-antarctic-rift-implications-volcanic-evolution.html#jCp>

Caitlin Gionfriddo, Science on ice: microbes and mercury in Antarctica, , Nature Research Microbiology, 8-2-2016, <https://naturemicrobiologycommunity.nature.com/users/18594-caitlin-gionfriddo/posts/11053-science-on-ice-microbes-and-mercury-in-antarctica>

Sebastian Goller, Antarctic subglacial hydrology - interactions of subglacial lakes, basal water flow and ice dynamics, 2014, <http://elib.suub.uni-bremen.de/peid=D00103852>

Serge Guillas, Simon Day, and B. McGuire, Statistical analysis of the El Niño–Southern Oscillation and sea-floor seismicity in the eastern tropical Pacific, 28 May 2010, <https://doi.org/10.1098/rsta.2010.0044>

Mark Hannington, P. Stoffers, I. C. Wright, and P. Herzig, Study shows that hydrothermal vents release mercury, Geological Survey of Canada, Ottawa, in the New Scientist, 10-6-1999, and Geology, vol 27, p 931, https://www.researchgate.net/publication/249520559_Elemental_mercury_at_submarine_hydrothermal_vents_in_the_Bay_of_Plenty_Taupo_volcanic_zone_New_Zealand

Lauren Hinkel, Southern Ocean cooling in a warming world, *Oceans at MIT* June 24, 2016, <http://news.mit.edu/2016/southern-ocean-cooling-in-a-warming-world-0624>

Ice Age Now, Underwater volcanic range discovered off coast of Norway, 4 August 2013, <https://www.iceagenow.info/12218/>

Akihiko Hokugo, Science Direct, Mechanism of Tsunami Fires after the Great East Japan Earthquake 2011 and Evacuation from the Tsunami Fires, *Volume 62*, 2013, Pages 140-153, <https://doi.org/10.1016/j.proeng.2013.08.051>

Barbel Hönisch, Andy Ridgwell, Naiela Schmidt, Ellen Thomas, Samantha Gibbs, Appy Sluijs, Richard Zeebe, Lee Kump, Rowan Martindale, Sarah Greene, Wolfgang Kiessling, Justin Ries, Jame Zachos, Dana Royer, Stephen Barker, Thomas Marchitto Jr., Ryan Moyer, Carles Pelejero, Patrizia Ziveri, Gavin Foster, and Branwen Williams, The Geological Record of Ocean Acidification, *Science* 02 Mar 2012: Vol. 335, Issue 6072, pp. 1058-1063, [DOI: 10.1126/science.1208277](https://doi.org/10.1126/science.1208277)

Evgenia Ilyinskaya, Stephen Mobbs, Ralph Burton, Mike Burton, Federica Pardini, Melissa Pfeffer, Ruth Purvis, James Lee, Stephane Bauguitte, Barbara Brooks, Ioana Colfescu, Gudrun Petersen, Axel Wellpot, and Baldur Bergsson, Globally Significant CO₂ Emissions From Katla, a Subglacial Volcano in Iceland, 17 Geophysical Research Letters, September 2018, <https://doi.org/10.1029/2018GL079096>

Erik Irvins, Ramp-up in Antarctic Ice Loss Speeds Sea Level Rise, NASA Earth Data Online, Jet Propulsion Laboratory. <https://sealevel.nasa.gov/news/129/ramp-up-in-antarctic-ice-loss-speeds-sea-level-rise> and https://www.nature.com/articles/s41586-018-0179-y.epdf?author_access_token=G6bM-sEvNrsr_d3FPj8qjtRgNOjAjWel9jnR3ZoTvOPBEKqWHTwARlrR4OxoHFdEh63arkDNI_bORoXuP_CQqP5K8MYc-mJnNFT_QmTd-WnNN5Mp3ZqXQU1Cq6c0OT0JzMpvEGDRBCqgg_mMZ20Fg%3D%3D

Nels Iverson, Ross Lieb-Lappen, Neila Dunbar, Rachel Obbard, Ellen Kim, and Ellyn Golden, The first physical evidence of subglacial volcanism under the West Antarctic Ice Sheet *Scientific Reports* volume 7, Article number: 11457 (2017), <https://www.nature.com/articles/s41598-017-11515-3>

O. Jagoutz, Study: Ancient tectonic activity was trigger for ice ages, MIT News, 18 April 2016, <http://news.mit.edu/2016/ancient-tectonic-activity-was-trigger-for-ice-ages-0418>

Stewart Jamieson, Neil Ross, Jamin Greenbaum, Duncan Young, Allan Aitken, Jason Roberts, Donald Blankenship, Sun Bo, and Martin Siegert, An extensive subglacial lake and canyon system in Princess Elizabeth Land, East 1 February 2016, *Antarctica Geology* (2016) 44 (2): 87-90, <https://doi.org/10.1130/G37220.1>

Hugh Jenkyns, Geochemistry of oceanic anoxic events, 9 March 2010, <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2009GC002788>

Dougal Jerram, Mike Widdowson, Paul B. Wignall, Yadong Sun, Xulong Lai, David P.G. Bond, Trond H. Torsvik, Submarine palaeoenvironments during Emeishan flood basalt volcanism, SW China: Implications for plume–lithosphere interaction during the Capitanian, Middle Permian ('end Guadalupian') extinction event <https://www.sciencedirect.com/science/article/pii/S0031018215003065>

Joel Johnson, Jurgen Mienert, Andreia Plaza-Faverola, Sunil Vadakkepuliambatta, Joel Knies, Stefan Bunz, Karin Andreassen, and Benedicte Ferre, Abiotic methane from ultraslow-spreading ridges can charge Arctic gas hydrates, *Geology* (2015) 43 (5): 371-374, 1 May 2015, <https://doi.org/10.1130/G36440.1>

T. A. Jordan, C. Martin, F. Ferracoli, M. Matsuoka, H. Corr, R. Forsberg, A. Olesen, and M. Siegert, Anomalously high geothermal flux near the South Pole, *Scientific Reports* volume 8, Article number: 16785 (2018), 14 November 2018, <https://www.nature.com/articles/s41598-018-35182-0>

J. E. Kamis, Plate Climatology Theory, retired geologist, 7 October 2014, <https://static1.squarespace.com/static/55315cdae4b03d5a7f6f23e1/t/5c363e5e40ec9a23cdde790c/1547058785801/PlateClimatologyTheory+7+October+2014.pdf>

J. E. Kamis, El Nino's Generated by Geological Heat Flow Not Global Warming, [plateclimatology.com](http://www.plateclimatology.com/el-nios-generated-by-geological-heat-flow-not-global-warming/), 12 January 2015, <http://www.plateclimatology.com/el-nios-generated-by-geological-heat-flow-not-global-warming/>

J. E. Kamis, How coral reefs Recover from Geologically Induced Deep Ocean "Forest Fires", [plateclimatology.com](http://www.plateclimatology.com/how-coral-reefs-recover-from-geologically-induced-deep-ocean-forest-fires/), 18 January 2015, <http://www.plateclimatology.com/how-coral-reefs-recover-from-geologically-induced-deep-ocean-forest-fires/>

J. E. Kamis, Fire and Ice, [plateclimatology.com](http://www.plateclimatology.com), 29 January 2015.

J. E. Kamis, Gulf Stream Shut Down Caused by Geological Heat flow, [plateclimatology.com](http://www.plateclimatology.com/gulf-stream-shut-down-caused-by-geological-heat-flow/), 16 March 2015, <http://www.plateclimatology.com/gulf-stream-shut-down-caused-by-geological-heat-flow/>

J. E. Kamis, How Megaplumes Influence the Climate Much More Than Expected, 9 April 2015, <http://www.plateclimatology.com/research2image/>

J. E. Kamis, Deep Ocean Rock Layer Mega-Fluid Flow Systems, 29 October 2015, [plateclimatology.com](http://www.plateclimatology.com)

J. E. Kamis and P. E. Kamis, Plate Climatology Theory, American Meteorological Society Conference, 13 January 2016, Poster Session, <https://ams.confex.com/ams/96Annual/webprogram/Paper290033.html>

J. E. Kamis, Heat from Deep Ocean Fault Punches Hole in Arctic Ice Sheet, [plateclimatology.com](http://www.plateclimatology.com), 6 November 2016 <http://www.plateclimatology.com/heat-from-deep-ocean-fault-punches-hole-in-arctic-ice-sheet/>

J. E. Kamis, Researchers discover volcanic activity is changing Antarctica's geology, [plateclimatology.com](http://www.plateclimatology.com), 6 February 2016. <http://www.plateclimatology.com/researchers-discover-volcanic-activity-is-changing-antarcticas-geology/>

J. E. Kamis, Super-Strong El Nino's are Nature's Underwater Forest Fires, [plateclimatology.com](http://www.plateclimatology.com), 15 April 2016, <http://www.plateclimatology.com/superstrong-el-nios-are-natures-underwater-forest-fires/>

J. E. Kamis, Active Volcanoes, Geological Forces Melting Antarctica's Nansen Glacier – Not Global Warming, Retired Geologist, 20 April 2016, [plateclimatology.com](http://www.plateclimatology.com), <http://www.plateclimatology.com/active-volcanoes-geological-forces-melting-antarcticas-nansen-glacier-not-global-warming/>

J. E. Kamis, Further Proof El Nino's are Fueled by Deep-Sea Geological Heat Flow, [plateclimatology.com](http://www.plateclimatology.com), 27 January 2017.

J. E. Kamis, In latest West Antarctic glacial melt scare influence of geological heat flow ignored, 9 October 2017, <http://www.plateclimatology.com/in-latest-west-antarctic-glacial-melt-scare-influence-of-geological-heat-flow-ignored/>

J. E. Kamis, NASA Study Shows Additional Geological Forces Behind West Antarctic Ice Melt, [plateclimatology.com](http://www.plateclimatology.com), 10 November 2017, <http://www.plateclimatology.com/nasa-study-shows-additional-geologic-forces-behind-west-antarctic-ice-melt/>

J. E. Kamis, 11 January 2018, How Major Oceanic Continental Fault Boundaries Act to Control Much of Earth's Climate, [plateclimatology.com](http://www.plateclimatology.com), <http://www.plateclimatology.com/how-major-oceanic-and-continental-fault-boundaries-act-to-control-much-of-earths-climate/>

J. E. Kamis, Mounting geological Evidence Shows Ice Sheet Melting Natural, [plateclimatology.com](http://www.plateclimatology.com), 18 January 2018, <http://www.plateclimatology.com/mounting-geological-evidence-shows-ice-sheet-melting-natural/>

J. E. Kamis, Largest High-Arctic Lake Melting from Geothermal Heat, Not Global Warming, [plateclimatology.com](http://www.plateclimatology.com), 16 April 2018.

J. E. Kamis, West Antarctic Volcano and Fault Belt Part of Pacific Ring of Fire, [plateclimatology.com](http://www.plateclimatology.com), 7 May 2018, <http://www.plateclimatology.com/west-antarctic-volcano-and-fault-belt-part-of-pacific-ring-of-fire/>

J. E. Kamis, Three New Studies Confirm Volcanism is Melting West Antarctic Glaciers, Not Global Warming, [plateclimatology.com](http://www.plateclimatology.com), 29 June 2018.

J. E. Kamis, Geological Activity, Not "Atlantification", Altering Arctic Ocean Temps and Salinity, plateclimatology.com, 16 July 2018.

Gerta Keller, Paula Mateo, Jahnavi Punekar, Hassan Khozyem, Brian Gertsch, Jorge Spangenberg, Andre Bitchong, nd Thierry Adatte, Environmental changes during the Cretaceous-Paleogene mass extinction and Paleocene-Eocene Thermal Maximum: Implications for the Anthropocene, *Gondwana Research* Volume 56, April 2018, Pages 69-89, <https://www.sciencedirect.com/science/article/pii/S1342937X17303702>

Jessica Klar, Rachael James, Dakota Gibbs, Alastair Lough, Ian Parkinson, Andrew Milton, Jeffrey Hawkes, and Douglas Connelly, Isotopic signature of dissolved iron delivered to the Southern Ocean from hydrothermal vents in the East Scotia Sea, *GSA Data Repository* 2017101, 2017, <http://www.geosociety.org/datarepository/2017/2017101.pdf>

G. P. Klinkhammer, C. S. Chin, R. A. Keller, A. Dahmann, H. Sahling, G. Sarthou, S. Petersen, F. Smith, and C. Wilson, Discovery of new hydrothermal vent sites in Bransfield Strait, Antarctica, *Science Direct*, 15 December 2001, [https://doi.org/10.1016/S0012-821X\(01\)00536-2](https://doi.org/10.1016/S0012-821X(01)00536-2)

Tormod Kvaerna, Seismology in the Svalbard Region, [TormodKvaernaSeismology%20\(6\)](#)

Kaoru Kubota, Yusuke Yokoyama, Tsuyoshi Ishikawa, Stephen Obrochita, and Atsushi Suzuki, Larger CO₂ source at the equatorial Pacific during the last deglaciation, *Science Reports* 4, article number:5261(2014), 11 June 2014, <https://www.nature.com/articles/srep05261#f1>

C. Läderach, V. Schlindwein, H. Schenke, and W. Jokat, Seismicity and active tectonic processes in the ultra-slow spreading Lena Trough, Arctic Ocean, *Geophysical Journal International*, Volume 184, Issue 3, 1 March 2011, Pages 1354–1370, <https://doi.org/10.1111/j.1365-246X.2010.04926.x>, 01 March 2011

Igor Lehnerr, Vincent St. Louis, Martin Sharp, Alex Gardner, John Smol, Sherry Schiff, Derek Muir, Colleen Mortimer, Neil Michelutti, Charles Tamoochai, Kyra, St. Pierre, Craig Emmerton, Johan Wiklund, Guiter Kock, Scott Lamoureux, and Charles Talbot, The World's largest High Arctic lake responds to climate change, *Nature Communications*, 9 article number:1290(2018), 29 March 2018, <https://www.nature.com/articles/s41467-018-035685-z>

Brice Loose, Alberto Garabato, Peter Schlosser, William Jenkins, David Vaughan, and Karen Heywood, Evidence of an active volcanic heat source beneath the Pine Island Glacier, *Nature Communications*, volume 9, Article number: 2431, 22 June 2018, https://www.nature.com/articles/s41467-018-04421-3?error=cookies_not_supported&code=2cdd024f-7455-4480-a95e-42e71144ed07

Amanda Lough, Douglas Wiens, C. Barcheck, Sridhar Anandkrishnan, Robert Aster, Donald Blankenship, Audrey Huerta, Andrew Nyblade, Duncan Young, and Terry Wilson, Seismic Detection of an active subglacial magmatic complex Marie Byrd Land Antarctica, *Nature Geoscience*, 17 November 2013, <https://www.nature.com/articles/ngeo1992> and https://www.nature.com/articles/ngeo1992.epdf?referrer_access_token=wLfsiQncvIR_Y5fEjzFs9RgN0jAjWeI9jnR3ZoTv0MAhwOSOkGjC4BaxrCpS698awgqJJUwuGe4iMiaowLzWqT_ndf_1oUtwrNDRHikR4K8Qf8osPnruPtloekPvU160BkT627zyyGBYBcTp9LdNWbQ44gvi68_Ls-OECQA5xLmM1Tiall5iG04oZRyxk84&tracking_referrer=news.nationalgeographic.com&error=cookies_not_supported&code=4a63b48f-eb47-40c8-bb47-1367cfee6207

Jennifer MacKinnon, Oceanographers Find Clues behind Arctic's Fourth-Lowest Sea-Ice Minimum, Scripps Institution of Oceanography, 16 September 2015, <https://scripps.ucsd.edu/news/oceanographers-find-clues-behind-arctics-fourth-lowest-sea-ice-minimum>

Adolfo Maestro, Paleomagnetic studies in Deception Island and Port Foster, Antarctica, Geomagdec: PI:(2008-2009), Volcano Discovery. https://www.volcanodiscovery.com/deception_island.html).

Ines Martins, Valentina Costa, Filipe Porteiro. and Ana Colaco, Mercury concentrations in fish species caught at Mid-Atlantic Ridge hydrothermal vent fields, I., Marine Ecology Progress Series, vol. 320: 253–258, 2006, https://www.researchgate.net/publication/229516990_Mercury_concentrations_in_fish_species_caught_at_Mid-Atlantic_Ridge_hydrothermal_vent_fields

Martos, Y., et. al. Thermal Scar From Dinosaur Era Still Warming Base of Greenland Ice Sheet, New Findings. NASA / BAS. <https://www.bas.ac.uk/media-post/thermal-scar-from-dinosaur-era-still-warming-base-of-greenland-ice-sheet-new-findings-show/>, 1 August 2018

Yamina Martos and Tom Jordan, New Antarctica heat map reveals sub-ice hotspots, British Antarctic Survey, Press Release, 13 November 2017, <https://www.bas.ac.uk/media-post/new-antarctic-heat-map-reveals-sub-ice-hotspots/>

Susan Mau, Miriam Romer, Marta Torre, and Ineborg Bussmann, Widespread methane seepage along the continental margin off Svalbard - from Bjørnøya to Kongsfjorden, *Scientific Reports* **volume7**, Article number: 42997 (2017), 23 February 2017, https://www.researchgate.net/publication/313962145_Widespread_methane_seepage_along_the_continental_margin_off_Svalbard_-_from_Bjornoya_to_Kongsfjorden

Joseph McConnell, Andrea Burke, Nelia Dunbar, Peter Kohler, Jennie Thomas, Monica Arienzo, Nathan Chellman, Olivia Maselli, Michael Sigl, Jess Adkins, Danial Baggenstos, John Burkhart, Edward Brook, Chisto Buizert, Jihong Cole-Dai, T. J. Fudge, Gregor Knorr, Hans Graf, Mackenzie Grieman, Nels Iverson, Kenneth McGwire, Robert Mulvaney, Guillaume Paris, Rachael Rhodes, Eric Saltzman, Jeffrey Severinghaus, Jorgen Steffensen, Kendrick Taylor, and Gisela Winckler, Synchronous volcanic eruptions and abrupt climate change ~17.7 ka plausibly linked by stratospheric ozone depletion, PNAS, 5 September 2017, <https://doi.org/10.1073/pnas.1705595114>

Malcolm McMillan, Anrew Shepherd, Noel Gourmelen, Amber Ridout, Thomas Flament, Anna Hogg, Lin Gilbert, Toby Benham, Michiel Van Den Broeke, Julian Dowdeswell, Xavier Fettweis, Brice Noel, and Tazio Strozzi, Rapid dynamic activation of a marine-based Arctic ice cap, Geophysical Research Letters, Volume 41, issue 2, 2 December 2014, <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1002/2014GL062255>

M. A. Martínez-Botí, G. Marino, G. L. Foster, P. Ziveri, M. J. Henehan, J. W. B. Rae, P. G. Mortyn, and D. Vance, Boron isotope evidence for oceanic carbon dioxide leakage during the last deglaciation, *Nature* volume 518, pages 219–222, 12 February 2015, <https://www.nature.com/articles/nature14155>.

Jurgen Mienert, New source of methane discovered in the Arctic Ocean, April 14, 2015, University of Tromso, Physics.org, <https://phys.org/news/2015-04-source-methane-arctic-ocean.html>

MIT, Volcanoes contribute to recent warming "hiatus", MIT News, 23 February 2014, <http://news.mit.edu/2014/study-volcanoes-contribute-to-recent-warming-hiatus-0223>

MIT, Small Volcanic Eruptions could be Slowing Global Warming, EAPS, 20 November 2014,
<https://eapsweb.mit.edu/news/2014/small-volcanic-eruptions-could-be-slowing-global-warming>

R. D. Muench, J. H. Morison, L. Padman, D. Martinson, P. Schlosser, B. Huber, and R. Hohmann, Maude Rise Revisited, *Journal of Geophysical Research-Oceans*, volume 106, p 2423-2440, 15 February 2001,
<https://www.ldeo.columbia.edu/node/10200>

Francesco Muschitiello, Francesco Pausata, James Lea, Douglas Mair, and Barbara Wohlfarth, Enhanced ice sheet melting driven by volcanic eruptions during the last deglaciation, *Nature Communications*, volume 8, Article number: 1020 (2017), 24 October 2017, https://www.nature.com/articles/s41467-017-01273-1?error=cookies_not_supported&code=04006c07-6911-463c-843f-7e3b05db3e4a

NASA, Antarctic Temperature Trend 1982-2004, NASA Earth Observatory News, 2006,
<https://earthobservatory.nasa.gov/images/6502/antarctic-temperature-trend-1982-2004>

NASA, NASA Satellite Detects Red Glow to Map Global Ocean Plant Health, 28 June 2009,
https://www.nasa.gov/topics/earth/features/modis_fluorescence.html

NASA, Hot News from the Antarctic Underground, *NASA Climate News*, 7 November 2017,
<https://www.nasa.gov/feature/jpl/hot-news-from-the-antarctic-underground>

Nat Geo, Ocean Acidification, National Geographic, 27 April 2017,
<https://www.nationalgeographic.com/environment/oceans/critical-issues-ocean-acidification/>

Julien Nicolas and David Bromwich, New Reconstruction of Antarctic Near-Surface Temperatures: Multidecadal Trends and Reliability of Global Re-Analyses, *AMS Journals Online*,
<https://journals.ametsoc.org/doi/10.1175/JCLI-D-13-00733.1>, <https://journals.ametsoc.org/doi/10.1175/JCLI-D-13-00733.1>

Grace Nield, Valentina Barletta, Andrea Bordoni, Matt King, Pippa Whitehouse, Peter Clarke, Eugene Domack, Ted Scambos, and Etienne Berthie, Rapid bedrock uplift in the Antarctic Peninsula explained by viscoelastic response to recent ice unloading, *Earth and Planetary Science Letters*, 5 May 2014,
<https://www.sciencedirect.com/science/article/pii/S0012821X14002519>

Bernt Olsen, Christofer Troedsson, Kenan Hadziavdic, Rolf-Birger Pedersen, and Hans Rapp, The influence of vent systems on pelagic eukaryotic micro-organism composition in the Nordic Seas, *Polar Biology*, Volume 38, Issue 4, pp 547–558, 25 November 2014, <https://link.springer.com/article/10.1007/s00300-014-1621-8>
Christopher Omelon, Wayne Pollard, and Dale Andersen, A geochemical evaluation of perennial spring activity and associated mineral precipitates at Expedition Fjord, Axel Heiberg Island, Canadian High Arctic, *Applied Geochemistry*, V 21 Issue 1, January 2006,
<https://www.sciencedirect.com/science/article/pii/S088329270500171X>

Oregon State University, South Sandwich Islands, southern Atlantic Ocean,
http://volcano.oregonstate.edu/vwdocs/volc_images/south_america/south_sandwich_islands.html

Ayako Abe-Ouchi, Fuyuki Saito, Kenji Kawamura, Maureen E. Raymo, Jun'ichi Okuno, Kunio Takahashi, and Heinz Blatter, Insolation-driven 100,000-year glacial cycles and hysteresis of ice-sheet volume, *Nature* **volume 500**, pages 190–193, 8 August 2013, <https://www.nature.com/articles/nature12374#article-info>

K. Pipejohn, W. von Gosen, F. Tessensohn, and Kerstin Saalmann, Ellesmerian fold-and thrust belt(northeast Ellesmere Island, Nunavut) and its Eureka overprint, Bulletin of the Geological Survey of Canada, 592 chapter 11, January 2008, https://www.researchgate.net/publication/262323657_Ellesmerian_fold-and-thrust_belt_northeast_Ellesmere_Island_Nunavut_and_its_Eureka_overprint

Rolf Pedersen and Terje Bjerkgard, Sea-Floor Massive Sulphides in Arctic Waters, Chapter 5, http://www.ngu.no/upload/Aktuelt/CircumArctic/5_SMS.pdf

W. H. Pollard, C. Omelon, and C. McKay, Geomorphic and Hydrologic Characteristics of Perennial Springs on Axel Heiberg Island, Canadian High Arctic, 1998, Seventh Annual Conference, Number 55, <https://www.tib.eu/en/search/id/BLCP%3ACN035207089/Geomorphic-and-hydrologic-characteristics-of-perennial/>

Igor Polyakov, Andrey Pnyushkov, Matthew Alkire, Igor Ashik, Till Baumann, Eddy Carmack, Ilonna Goszczko, John Guthrie, Vladimir Ivanov, Torsten Kanzow, Richard KrisField, Ronald Kwok, Arild Sundfjord, James Morison, Robert Rember, Alexander Yulin, Greater role for Atlantic inflows on sea-ice loss in the Eurasian Basin of the Arctic Ocean, *Science* 21 Apr 2017: Vol. 356, Issue 6335, pp. 285-291, DOI: 10.1126/science.aai8204, <http://science.sciencemag.org/content/356/6335/285>

J. Rey, L. Somoza, and J. Martinez-Frias, Tectonic, volcanic, and hydrothermal event sequence on Deception Island, Antarctica, May 23, 1994, in *Geo-Marine Letters* 1995:15 1-8, Hydrodec: Tectonic, volcanic and hydrothermal events in Deception Island (Antarctica). PI: Dr. Luis Somoza Losada (2000-2001).

Soroush Rezvanbehbahani, Machine learning predicts new details of geothermal heat flux beneath the Greenland Ice Sheet University of Kansas Online News, <http://today.ku.edu/2018/01/08/machine-learning-predicts-new-details-geothermal-heat-flux-beneath-greenland-ice-sheet>, 1 October 2018.

Alex Rogers, Paul Tyler, Douglas Connelly, Jon Copley, Rachael James, Robert Larter, Katrin Linse, Rachael Mills, Alfredo Garabato, Richard Pancost, David Pearce, Nicholas Polunin, Christopher German, Timothy Shank, Philipp Boersch-Supan, Belinda Alker, Alfred Aquilina, Sarah Bennett, Andrew Clarke, Robert Dinley, Alastair Graham, Darryl Green, Jeffrey Hawkes, Laura Hepburn, Ana Hilario, Veerle Huvenne, Leigh Marsh, Eva Ramirez-Llodra, William Reid, Christopher Roterman, Christopher Sweeting, Sven Thatje, and Katrin Zwirgmaier, The Discovery of New Deep-Sea Hydrothermal Vent Communities in the Southern Ocean and Implications for Biogeography, PLOS, January 3, 2012, <https://doi.org/10.1371/journal.pbio.1001234>

Irna Rogozhina, Alexey Petrunin, Alan Vaughan, Bernhard Steinberger, Jesse Johnson, Mikhail Kaban, Reinhard Calov, Florian Rickers, Maik Thomas, and Ivan Koulakov, Melting at the base of the Greenland ice sheet explained by Iceland hotspot history, *Nature Geoscience* volume9, pages366–369 (2016), 04 April 2016, <https://www.nature.com/articles/ngeo2689>

Anja Rutishauser, Donald Blankenship, Martin Sharp, Mark Skidmore, Jamin Greenbaum, Cyril Grima, Dustin Schroeder, Julian Dowdeswell, and Duncan Young, Discovery of a hypersaline subglacial lake complex beneath Devon Ice Cap, Canadian Arctic, *Science Advances* 11 Apr 2018: Vol. 4, no. 4, eaar4353, DOI: 10.1126/sciadv.aar4353, <http://advances.sciencemag.org/content/4/4/eaar4353.full>

Soren Rysgaard, Jorgen Bendtsen, John Mortesen, and Mikael Sejr, High geothermal heat flux in close proximity to the Northeast Greenland Ice Stream, *Scientific Reports* volume 8, Article number: 1344 (2018), 22 January 2018, <https://www.nature.com/articles/s41598-018-19244-x#Abs1>

Mak Saito, Abigail Noble, Alessandro Tagliabue, Tyler Goepfert, Carl Lamboug, and William Jenkins, Slow-spreading submarine ridges in the South Atlantic as a significant oceanic iron source, *Nature Geoscience*, 18 August 2013, DOI:10.1038/NGEO1893.

Haley Sapers, Jennifer Ronholm, Isabelle Raymond-Bouchard, Raven Comrey, Gordon Isinski, and Lyle Whyte, Biological Characterization of Microenvironments in a Hyper saline Cold Springs Mars Analog, *Frontiers in Microbiology*, doi: [10.3389/fmicb.2017.02527](https://doi.org/10.3389/fmicb.2017.02527), 22 December 2017, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5744183/>

Eric Schandl, Michael Gorton, and J. Wicks, Geochemistry of basalts from ODP Hole 113-690C, PANGAEA, 1990, DOI:10.1594/PANGAEA.753879, oai:pangaea.de, doi:10.1594/PANGAEA.753879, <https://core.ac.uk/display/57749100>

Vera Schlindwein, Christian Muller, and Wilfried Jokat, Seismoacoustic evidence for volcanic activity on the ultraslow-spreading Gakkel Ridge, Arctic Ocean, *Geophysical Research Letters*, 29 September 2005, <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2005GL023767>

Vera Schlindwein and F. Schmid, Mid-ocean-ridge seismicity reveals extreme types of ocean lithosphere, *Nature* volume 535, pages 276–279, 14 July 2016, <https://www.ncbi.nlm.nih.gov/pubmed/27362231>

Schmidtko, Sunke, et. al., Multidecadal warming of Antarctic waters, , *Science* 05 Dec 2014:Vol. 346, Issue 6214, pp. 1227-1231, DOI: 10.1126/science.1256117, <http://science.sciencemag.org/content/346/6214/1227>

Dustin Schroeder, Donald Blankenship, Duncan Young, and Enrica Quartini, Evidence for elevated and spatially variable geothermal flux beneath the West Antarctic Ice Sheet, *Proceedings of the National Academy of Sciences*, June 2014, DOI: 10.1073/pnas.1405184111. <https://www.pnas.org/content/111/25/9070>

Al. Schreidera, A. Schreider, J. Galindo-Zaldivar, A. Maldonado, L Gamboa, Y. Martos, and E. Evsenko, Structure of the Bransfield Strait Crust, *Marine Geology*, November 19, 2013, <https://link.springer.com/article/10.1134/S0001437014060101>

M. Scott, 2018 Arctic Report Card: Multi-year stretch of record and near-record warmth unlike any period on record, NOAA, 11 December 2018

Sergio Sejas, Unmasking the negative greenhouse effect over the Antarctic Plateau, July 2018, DOI: 10.1038/s41612-018-0031-y

Helene Seroussi, Erik Ivins, Douglas Wiens, and Johannes Bondzio, Influence of a West Antarctic mantle plume on ice sheet basal conditions, 01 August 2017, <https://doi.org/10.1002/2017JB014423>

Martin Siegert, Neil Ross, and Anne Le Brocq, Recent advances in understanding Antarctic subglacial lakes and hydrology, *Philos Trans A Math Phys Eng Sci.*, 2016 Jan 28; 374(2059): 20140306., [10.1098/rsta.2014.0306](https://doi.org/10.1098/rsta.2014.0306)

Smithsonian Institution, Global Volcanism Program, Erebus, <https://volcano.si.edu/>

Smithsonian Institution, GVP, <https://volcano.si.edu/volcano.cfm?vn=234011>

Jørgen Peder Steffensen, Katrine K. Andersen, Matthias Bigler, Henrik B. Clausen, Dorthe Dahl-Jensen1, Hubertus Fischer, Kumiko Goto-Azuma, Margareta Hansson, Sigfús J. Johnsen, Jean Jouzel, Valérie Masson-Delmotte, Trevor Popp, Sune O. Rasmussen, Regine Röthlisberger, Urs Ruth, Bernhard Stauffer, Marie-Louise

Siggaard-Andersen, Árný E. Sveinbjörnsdóttir, Anders Svensso, and James W. C. White, High-Resolution Greenland Ice Core Data Show Abrupt Climate Change Happens in Few Years, *Science* 01 Aug 2008: Vol. 321, Issue 5889, pp. 680-684, DOI: 10.1126/science. 1157707
<http://science.sciencemag.org/content/321/5889/680.full>

Eric Steig, David Schneider, Scott Rutherford, Michael Mann, Josefino Comiso, and Drew Shindell, Warming of the Antarctic ice-sheet surface since the 1957 International Geophysical Year, *Nature* volume 457, pages 459–462 (22 January 2009), <https://www.nature.com/articles/nature07669>

Eric Steig, NASA Study: Mass Gains of Antarctic Ice Sheet Greater than Losses, NASA latest News, 30 October 2015.

Malte Stuecker, Cecilia Bitz, Kyle Armour, Cristian Proistosescu, Srah Kang, Shang-Ping Xie, Doyeon Kim, Shayne McGregor, Wenjun Zhang, Sen Zhao, Wenju Cai, Yue Dong, and Fei-Fei Jin, Polar amplification dominated by local forcing and feedbacks, *Nature Climate Change* (2018), <https://www.nature.com/articles/s41558-018-0339-y>

S Swart, EC Campbell, CH Heuze, K Johnson, JL Lieser, R Masson, M Mazloff, M Meredith, P Reid, J-B Sallee and S Stammerjohn, Return of the Maude Rise Polynya: Climate Litmus or Sea Ice anomaly, *Bulletin of the American Meteorological Society*, vol. 99, no. 8 , S188-S189 , 2017, <https://eprints.utas.edu.au/27884/>.

Theodore Them II, Benjamin Gill, Andrew Caruthers, Angela Gerhardt, Darren Grocke, Timothy Lyons, Selva Marroquin, Sune Nielsen, Joao Alexandre, and Jeremy Owens, Thallium isotopes reveal protracted anoxia during the Toarcian (Early Jurassic) associated with volcanism, carbon burial, and mass extinction, *PNAS* June 26, 2018 115 (26) 6596-6601; published ahead of print June 11, 2018 <https://doi.org/10.1073/pnas.1803478115>

Timmermans, M. L., Warming of the interior Arctic Ocean linked to sea ice losses at the basin margins, *Science Advances*, 29 Aug 2018:Vol. 4, no. 8, eaat6773, DOI: 10.1126/sciadv.aat6773

Maya Tolstoy, Seafloor Volcano Pulses May Alter Climate, Columbia University's Lamont-Doherty Earth Observatory 5 February 2015, <https://www.ldeo.columbia.edu/news-events/seafloor-volcano-pulses-may-alter-climate>.

Maya Tolstoy, Seafloor volcano pulses may alter climate: Strikingly regular patterns, from weeks to eons, *Science Daily*, 5 February 2015, <https://www.sciencedaily.com/releases/2015/02/150205142921.htm>

John Turner, Hua Lu, Ian white, John King, Tony Phillips, J. Hosking, Thomas Bracegirdle, Gareth Marshall, Robert Mulvaney, and Pranab Deb, Absence of 21st century warming on Antarctic Peninsula consistent with natural variability, *Nature* volume 535, pages 411–415, 21 July 2016.

University of Bergen, Seismicity of Norway and Surrounding Areas, 2013, http://nnsn.geo.uib.no/reports/2013/seismicity_2013.pdf

Costantino Vetriani, Yein Chew, Susan Miller, Jane Yagi, Jonna Coombs, Richard Lutz, and Tamar Barkay, Mercury Adaptation among Bacteria from a Deep-Sea Hydrothermal Vent, *Appl Environ Microbiol.* 2005 Jan; 71(1): 220–226, doi: 10.1128/AEM.71.1.220-226.2005, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC544242/>

Volcano Discovery, The South Shetland Islands Deception Island Caldera, https://www.volcanodiscovery.com/deception_island.html

Volcano Discovery, Mount Erebus, <https://www.volcanodiscovery.com/erebus.html>

Volcano Live, <http://volcanolive.com/mcdonaldisland.html>

Duncan Wingham, Martin Siegert, Andrew Shepherd, and Alan Muir, Rapid discharge connects Antarctic subglacial lakes, *Nature Letters*, Vol 440 | 20 April 2006 | doi:10.1038/nature04660, https://courses.eas.ualberta.ca/eas570/wingham_etal.nature06.pdf

Andrew Wright and Martin Siegert , [A fourth inventory of Antarctic subglacial lakes](#). *Antarctic Science* 24, 659-664, 2012

James Zachos, Ursula Rohl, Stephen Schellenberg, Appy Sluijs, David Hodell, Daniel Kelly, Ellen Thomas, Micah Nicolo, Isabella Raffi, Lucas Lourens, Heather McCarren, and Dick Kroon, Rapid Acidification of the Ocean During the Paleocene-Eocene Thermal Maximum, *Science*, 10 Jun 2005: Vol. 308, Issue 5728, pp. 1611-1615, DOI: 10.1126/science.1109004, <http://science.sciencemag.org/content/308/5728/1611>

D. Zandomenighi, R. Aster, P. Kyle, A. Barclay, J. Chaput, and H. Knox, Internal structure of Erebus volcano, Antarctica imaged by high-resolution active-source seismic tomography and coda interferometry, *Journal of Geophysical Research: solid earth*, vol. 118, 1067–1078, <https://doi.org/10.1002/jgrb.50073>, 13 January 2013, <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1002/jgrb.50073>

Xin-Yuan Zheng, Hugh Jenkyns, Andrew Gale, David Ward, and Gideon Henderson, Changing ocean circulation and hydrothermal inputs during Ocean Anoxic Event 2 (Cenomanian–Turonian): Evidence from Nd-isotopes in the European shelf sea, *Earth and Planetary Science Letters*, *Volume 375*, 1 August 2013, Pages 338-348, <https://doi.org/10.1016/j.epsl.2013.05.053>

Jay Zwally, NASA Study: Mass Gains of Antarctic Ice Sheet Greater than Losses, *NASA News*, 30 October 2015, <https://climate.nasa.gov/news/2361/study-mass-gains-of-antarctic-ice-sheet-greater-than-losses/>